

Digitized by the Internet Archive
in 2011 with funding from
LYRASIS members and Sloan Foundation

Bulletin 366

January, 1935

S
43
E22
no. 366

REPORT OF THE DIRECTOR

For the Year Ending

October 31, 1934



Connecticut
Agricultural Experiment Station
New Haven

CONNECTICUT AGRICULTURAL EXPERIMENT STATION

(As of October 31, 1934)

BOARD OF CONTROL

His Excellency, Governor Wilbur L. Cross, *ex-officio*, President

Elijah Rogers, <i>Vice-President</i>	Southington
William L. Slate, <i>Treasurer</i>	New Haven
Edward C. Schneider, <i>Secretary</i>	Middletown
Joseph W. Alsop	Avon
Charles G. Morris	Newtown
Albert B. Plant	Branford
Olcott F. King	South Windsor

STAFF

Administration.

WILLIAM L. SLATE, B.Sc., *Director*.
 MISS L. M. BRAUTLECHT, *Bookkeeper and Librarian*.
 MISS KATHERINE M. PALMER, B.LITT., *Editor*.
 G. E. GRAHAM, *In Charge of Buildings and Grounds*.

Analytical Chemistry.

E. M. BAILEY, PH.D., *Chemist in Charge*.
 C. E. SHEPARD
 OWEN L. NOLAN
 HARRY J. FISHER, PH.D. } *Assistant Chemists*.
 W. T. MATHIS
 DAVID C. WALDEN, B.S. }
 FRANK C. SHELDON, *Laboratory Assistant*.
 V. L. CHURCHILL, *Sampling Agent*.
 MRS. A. B. VOSBURGH, *Secretary*.

Biochemistry.

H. B. VICKERY, PH.D., *Biochemist in Charge*.
 LAFAYETTE B. MENDEL, PH.D., *Research Associate* (Yale University).
 GEORGE W. PUCHER, PH.D., *Assistant Biochemist*.

Botany.

G. P. CLINTON, Sc.D., *Botanist in Charge*.
 E. M. STODDARD, B.S., *Pomologist*.
 MISS FLORENCE A. McCORMICK, PH.D., *Pathologist*.
 A. A. DUNLAP, PH.D., *Assistant Mycologist*.
 A. D. McDONNELL, *General Assistant*.
 MRS. W. W. KELSEY, *Secretary*.

Entomology.

W. E. BRITTON, PH.D., D.Sc., *Entomologist in Charge, State Entomologist*.
 B. H. WALDEN, B.AGR.
 M. P. ZAPPE, B.S. } *Assistant Entomologists*.
 PHILIP GARMAN, PH.D.
 ROGER B. FRIEND, PH.D.
 NEELY TURNER, M.A. }
 JOHN T. ASHWORTH, *Deputy in Charge of Gipsy Moth Control*.
 R. C. BOTSFORD, *Deputy in Charge of Mosquito Elimination*.
 J. P. JOHNSON, B.S., *Deputy in Charge of Japanese Beetle Quarantine*.
 MISS HELEN A. HULSE } *Secretaries*.
 MISS BETTY SCOVILLE }

Forestry.

WALTER O. FILLEY, *Forester in Charge*.
 H. W. HICOCK, M.F., *Assistant Forester*.
 J. E. RILEY, JR., M.F., *In Charge of Blister Rust Control*.
 MISS PAULINE A. MERCHANT, *Secretary*.

Plant Breeding.

DONALD F. JONES, Sc.D., *Geneticist in Charge*.
 W. RALPH SINGLETON, Sc.D., *Assistant Geneticist*.
 LAWRENCE C. CURTIS, B.S., *Assistant*.
 MRS. GENEVIEVE B. DEWEY, A.B., *Secretary*.

Soils.

M. F. MORGAN, M.S., *Agronomist in Charge*.
 H. G. M. JACOBSON, M.S., *Assistant Agronomist*.
 HERBERT A. LUNT, PH.D., *Assistant in Forest Soils*.
 DWIGHT B. DOWNS, *General Assistant*.
 MISS GERALDINE EVERETT, *Secretary*.

Tobacco Substation at Windsor.

PAUL J. ANDERSON, PH.D., *Pathologist in Charge*.
 T. R. SWANECK, M.S., *Agronomist*.
 O. E. STREET, PH.D., *Plant Physiologist*.
 MISS DOROTHY LENARD, *Secretary*.


REPORT OF THE DIRECTOR

For the Year Ending

October 31, 1934



Connecticut
Agricultural Experiment Station
New Haven



STATION FIELD DAY, AUGUST 22, 1934

Members of the Station Board and Dr. Ezekiel, the speaker, are standing outside the big tent at the experiment farm at Mount Carmel. They are (left to right): A. B. Plant, Elijah Rogers, Director Slate, Governor Cross, Olcott King, Dr. Ezekiel, Charles G. Morris and E. C. Schneider.

REPORT OF THE DIRECTOR

For the Year Ending October 31, 1934

To the Board of Control of the Connecticut Agricultural Station:

WHATEVER may have been the experience in other groups, the past year has been one of cheerfulness and enthusiasm at the Station. In spite of reduced budgets and salaries, our work has gone well and quite smoothly. Perhaps this is the reflection of the better feeling generally prevalent among farmers; it may be due in part to the opportunity for driving certain slow-moving projects toward completion through the help of federal relief agencies. Whatever the cause, the effect has been to clear the skies and give us a renewed faith in the task of "putting science at work for agriculture".

The place of the Station in the affairs of the State has been re-emphasized, not only in the minds of the staff, but in the consciousness of the general public. Established primarily to serve agriculture, it has come to serve all citizens as a source of scientific information on a great variety of matters, both public and private. It has the advantage of maintaining a permanent body of scientists accumulating data on the problems of farmer and consumer over a long period of time. Thus in emergencies, no time is lost while new agencies or men familiarize themselves with a situation. The Station is ready to serve.

Recent illustrations are furnished in the cases of the Dutch elm disease, the soil testing service, the "X" disease of peaches and the expanded mosquito elimination project, discussed in more detail elsewhere in this report. Not that all questions can be answered when they arise. There is much yet to be learned about most problems, but we are in a strategic position. With no "program" to promote or carry out, with long experience in fact-finding, the Agricultural Station is unique among public institutions.

However, we may not rest complacently on past achievements or even in the present. As the *Field* of London stated recently: "Agriculture is changing so fast that experience of past methods is less useful than knowledge of new ones and the mental adaptability and courage necessary to try them. This alone goes a long way to justify expenditure on agricultural education and research".

A most interesting experience has been the Station's part in several federal relief projects. These have done double duty by speeding up the control of pests and by giving many of the unemployed productive work. At one time we were directing more than 2000 men. The largest group, more than 1000, was on mosquito elimination; about 200 helped in the Gipsy Moth control; Dutch elm disease, European pine shoot moth, and white pine blister rust accounted for the balance.

Field Days and Station Bulletins

Field days and bulletins are two of the means used by the Station to disseminate information. There were four field days in 1934: two at the farm at Mount Carmel and an early and a late vegetable field day at the Substation at Windsor. All of these were well attended, but the annual Station Field Day on August 22 was one of the best the Station has ever held. At the last minute Secretary of Agriculture, Henry A. Wallace, who was to address the meeting, was detained by the death of Speaker Rainey. Nevertheless about 1500 persons, farmers, and men and women representing every walk of life, inspected the exhibits and experimental plots and after lunch gathered in a big tent pitched on the alfalfa field. Director Slate and Governor Wilbur L. Cross, chairman of the Station Board, made short addresses. Then the Director introduced Dr. Mordecai Ezekiel, economic adviser to Secretary Wallace, who had been sent from Washington to represent the Secretary.

Dr. Ezekiel's topic was: The Agricultural New Deal—what it is; how it works and what it is attempting to do; and what it means to New England. His succinct explanation of each part was well received by the audience. In discussing the AAA program, he emphasized the point that the ultimate aim is not to reduce, but to increase, production. Quoting recent statistics, he showed that a good standard of living for everyone would demand such an increase. Meanwhile the AAA believes it is necessary for farmers to curtail their production just as manufacturers reduce their output to meet market demand. He said that New England had already felt the benefits of larger farm income in greater sales of manufactured goods to rural people in the West. Her tobacco farmers have been testing production control with good results, and the milk producers may later adopt a plan.

During the year many members of the staff have addressed groups of growers and of scientists telling about their work on some special problem or the advancement in one line of research. A list of publications for the year is appended to this report.

Noteworthy Station Publications

Among the station bulletins demanding special comment are the *Plant Pest Handbook, Part II, Diseases and Injuries*, by Dr. G. P. Clinton; and *Tobacco Culture in Connecticut* by Dr. P. J. Anderson, in charge of the Tobacco Substation at Windsor. These two books are valuable contributions to the world of science and to the farmer and gardener. Dr. Clinton has been Station Botanist for more than 30 years. His work has been recognized by national groups by his election as fellow of the

National Academy of Sciences and of the American Academy of Arts and Sciences.

The *Plant Pest Handbook* is the summation of his observations and studies over this long period in Connecticut. All diseases of plants that have come to his attention are listed, together with a brief discussion of the symptoms, cause, and, where possible, the control or cure. Together with *Part I, Insects*, published by Dr. W. E. Britton last year, this book forms a cyclopedia of plant pest information for Connecticut.

Dr. Anderson's bulletin is a history as well as a guide to tobacco growing in the State. After a brief review of tobacco culture from Indian days to the present, the author describes in detail the varieties of tobacco planted, the methods of cultivation from preparation of seedbeds to curing, and the insect and disease enemies of tobacco, with controls.

**Staff Members
Honored**

Several members of the staff have been honored by outside organizations in recognition of their scientific contributions. In June, Dr. Donald F. Jones was elected a fellow of the American Academy of Arts and Sciences.

At the request of Governor Cross, the Director has served since December, 1933, as Chairman of the State Planning Board. This has required spending considerable time in Hartford and a number of trips to Boston. However, the possibilities for public service of a peculiar type are almost unlimited and the experience has been very interesting. In accordance with the Governor's wishes, the Planning Board has kept its program to "fact-finding" projects rather than "planning". Because the function of the Station is primarily fact-finding, it would seem that our experience and point of view should be valuable.

SPECIAL INVESTIGATIONS

The Dutch Elm Disease in Connecticut

The departments of botany, forestry and entomology have been particularly concerned with the Dutch elm disease. While this disease is caused by a fungus, *Graphium ulmi*, and is therefore a botanical problem, it is carried from infected to healthy trees by certain beetles and so involves the entomologists. The fact that the fungus attacks one of New England's most valued shade trees makes it a vital concern of the forester also.

The Station Botanist is thoroughly familiar with *Graphium ulmi*. Ever since the outbreak of the disease in the Netherlands fifteen years ago, he has kept in close touch with scientific investigations, and has constantly watched for its possible appearance in Connecticut. Thus, when it was discovered in New York and New Jersey in 1933, the Station was prepared to meet the emergency.

Immediately the Botanist made a preliminary survey of elms in the State. No cases of *Graphium* were found at that time. Later in the autumn of 1933, a single case was located in Glenville, Connecticut, close to Westchester County, New York, where the disease is present.

During the winter, the Federal Government set up an office for Dutch elm disease control at Stamford and the Station coöperated with the activities of this agency. Together they supervised a crew of CWA workers in an intensive scouting campaign of southern Fairfield County. This resulted in the discovery of a second diseased tree.

**57 Diseased
Trees Found
In One Year**

After the CWA disbanded in May, a few experienced federal workers continued to scout in the section around Greenwich. Whenever an elm showed visible symptoms of disease—leaves wilting and turning yellow, and brown discoloration in the outer rings and under the bark of a cross-section—specimens were sent to the federal laboratory at Morristown, New Jersey, and to the station laboratory at New Haven. The field symptoms of *Graphium* so closely resemble other elm troubles that diagnosis can be made by laboratory culture only. Before the end of the summer, 56 specimens had been confirmed as *Graphium*. Diseased trees were cut down and burned.

In August and September the Station made a second, more intensive, state-wide survey of elm trees. Every town was visited, with the exception of that part of Fairfield County covered by the federal scouts. Of the many trees examined, 120 appeared to have symptoms of *Graphium* and samples were sent to the station laboratory. One tree only yielded a culture of *Graphium*. Further intensive search failed to locate any other cases in the region and the occurrence of this isolated diseased tree is yet a mystery.

The Botanist reports the significance of this one case: "It was found at Black Hall, Old Lyme, 50 miles from Fairfield County. The bark

contained the fruiting stage of the fungi more abundantly than any other tree in Connecticut to date. The European elm bark beetle, *Scolytus multistriatus*, known carrier of *Graphium*, was not present, nor was this

A New Carrier Found

insect found in the vicinity. Instead, the mature beetles and larvae of the native *Hylurgopinus rufipes* were present, as well as mites. These beetles, their larvae and the mites were found to be local carriers over the infested bark, since all of them were able to transfer the *Graphium* to media in Petri dishes."

The adult beetles also proved to be disease carriers. Placed in a test tube with twigs of healthy elm, they immediately bored into the bark. Later a culture from these twigs gave positive evidence of *Graphium*.

More than 50 dead or dying trees reported by the scouts were examined by the botanists. None of them contained *Graphium*, although a saprophytic *Graphium* did appear in some. So far the asco stage of *Graphium ulmi* has not been found in nature in this country. In a few cases the conidial stage has been produced by spraying pure culture of the spores on elm bark in moist chambers.

The scouts also made maps of the roads covered, indicating the condition of elms as good, fair and poor. These will be used in further surveying and scouting next spring. Trees reported to be in poor condition are being watched.

Committee on Elm Disease Formed

In October, interested citizens organized a Connecticut Committee on Dutch Elm Disease and the Station Forester was elected chairman. This local group co-operates with the New England Committee to urge federal action on the disease and to carry on a local educational campaign.

Acting together, New York, New Jersey and Connecticut appealed to the Federal Government to assign an emergency allotment of \$500,000 of PWA funds to finance immediate sanitation of infected areas. By sanitation is meant not only destruction of all diseased trees, but of elms, and parts of elms, dying from any cause. These measures carried out during this winter should reduce the number of beetles that hibernate under the bark of weak elms, thus eliminating possible sources of infection.

Governor Cross has taken an active interest. He has given all possible aid and support to both state and federal programs in their fight to save the elms of Connecticut.

A Comparison of the Weather Records at the Station Farm and the Weather Bureau Station in New Haven

The importance of accurate weather records to many fields of human endeavor, especially agriculture, is quite obvious. But in scientific investigations with plants and insects, their great value is often overlooked. The disturbing factor is the local variations that occur, particularly in a region with the location and topographic irregularities that are found in Connecticut. This State is bounded on the south by Long Island Sound and cut through the middle by the Connecticut River. Along the shore of the

sound and in the valley of this river the climate is somewhat milder than throughout the rest of the State. Two localities within a relatively short distance of each other may show considerable differences in weather conditions at any one time. This is particularly evident in temperature, in precipitation, and in wind velocity.

In 1931, we installed a fairly complete set of weather recording instruments on the experimental farm at Mount Carmel, Hamden. Prior to that time, dependence was placed on the records of the Weather Bureau at New Haven. A comparison of the weather records of the last three years, taken at the United States Weather Bureau Station at New Haven and at Mount Carmel, 7.5 miles north of the New Haven weather station and at a somewhat higher elevation, are very enlightening and demonstrate the need for many more weather stations in the State.

The instruments at the weather bureau station at New Haven are about 74 feet above sea level and in the center of the city, about one mile north of the shore of Long Island Sound. The instruments at the station farm are about 220 feet above sea level and 8.5 miles north of the sound. A low range of hills extends from a point about three miles northeast of the farm westward and then southward to terminate at West Rock, six miles south-southwest of the farm. The height of these hills varies from about 300 to 737 feet above sea level, the highest point, the Sleeping Giant or Mount Carmel, being about two miles north and slightly east of the farm. Directly north of the latter, at a distance of about one mile, the elevation is about 300 feet. Low land separates the farm from these hills to the north and west. The natural air drainage from the inland regions toward the sound is checked by this range of hills, and this probably accounts in part for lower winter temperatures which frequently occur on the northern side of the elevation.

In the following table, the mean monthly temperature and the monthly precipitation at Mount Carmel for the years 1932 to 1934 inclusive are compared with those of New Haven for the same years and with the New Haven normal means, based on records over a period from 1873 to date. The mean monthly temperature at Mount Carmel is usually between two and three degrees lower than that of New Haven, except for April and May, when it is between one and two degrees lower. During the winter months, the extremes of low temperature show a greater difference. During the period involved, the temperature at New Haven registered +10 degrees Fahrenheit or lower on 27 different days, whereas at Mount Carmel this occurred on 61 days. On 16 of these days the Mount Carmel temperature went to between 10 degrees and 16 degrees below that of New Haven. This is important in regard to injury to plants by cold and to the mortality of insect pests.

During the hottest days of summer, when the temperature is above 90 degrees Fahrenheit, the instruments at Mount Carmel usually register from one to four degrees above New Haven. The wind direction and velocity probably influence the extent to which the temperature at the two localities vary.

**Monthly Temperature and Precipitation at New Haven, Conn., and
Mount Carmel, Hamden, Conn., 1932 to 1934 Inclusive**

Year	Month	Mean temperature			Total precipitation		
		Mt. Carmel	New Haven	New Haven normal	Mt. Carmel	New Haven	New Haven normal
1932	Jan.	36.7°	39.6°	28.3°	4.95"	5.63"	3.98"
	Feb.	30.6	33.0	29.1	2.50	2.51	4.16
	Mar.	33.6	35.8	35.8	5.35	6.28	4.10
	Apr.	45.6	46.8	47.2	2.83	1.93	3.52
	May	58.0	59.4	57.9	2.00	2.64	3.69
	June	66.1	67.0	66.6	2.12	2.16	3.10
	July	70.7	72.5	71.6	1.77	2.79	4.32
	Aug.	70.4	72.8	70.3	4.40	3.88	4.26
	Sept.	62.8	65.1	63.6	3.65	3.55	3.54
	Oct.	52.9	56.1	53.7	5.28	5.51	3.66
	Nov.	39.1	41.7	42.0	6.72	6.45	3.35
	Dec.	32.7	36.1	32.5	2.43	2.25	4.01
Total					44.00"	45.58"	45.69"
1933	Jan.	34.2°	37.2°	28.2°	2.12"	1.73"	3.98"
	Feb.	29.6	32.4	29.0	5.00	4.15	4.00
	Mar.	34.7	36.7	35.8	6.30	6.60	4.10
	Apr.	46.1	47.4	47.2	4.98	4.64	3.52
	May	59.7	61.4	57.9	2.55	2.00	3.69
	June	67.2	69.2	66.6	2.70	2.26	3.10
	July	69.5	72.4	71.8	3.17	3.58	4.32
	Aug.	69.2	66.8	70.3	6.70	7.39	4.32
	Sept.	64.1	66.6	63.5	5.66	5.05	3.54
	Oct.	50.3	53.6	53.8	3.10	2.64	3.66
	Nov.	37.1	39.4	42.0	.77	.90	3.35
	Dec.	26.0	29.2	32.5	3.54	4.51	4.01
Total					46.59"	45.75"	45.59"
1934	Jan.	29.3°	31.6°	28.2°	4.02"	3.52"	3.98"
	Feb.	14.0	17.4	29.0	3.39	3.82	4.00
	Mar.	32.5	35.4	35.8	3.18	4.68	4.10
	Apr.	47.2	48.4	47.2	7.82	4.98	3.52
	May	59.2	60.4	57.9	6.66	5.23	3.69
	June	68.2	70.0	66.6	4.56	3.42	3.10
	July	72.3	74.8	71.8	2.34	3.04	4.32
	Aug.	65.7	68.6	70.3	2.19	2.45	4.26
	Sept.	64.8	66.5	63.5	11.07	8.76	3.54
	Oct.	48.8	51.6	53.8	2.72	2.51	3.66
	Nov.	44.2	46.7	42.0	4.35	3.28	3.35
	Dec.	28.7	31.2	32.5	4.99	3.34	4.01
Total					57.29"	49.03"	45.53"

In estimating the mean temperatures, the procedure differs at the two stations. Daily means are determined by averaging bi-hourly temperatures at Mount Carmel, and the monthly mean is the average of the daily means.

At New Haven the daily mean is half the sum of the maximum plus the minimum, and the monthly mean is half the sum of the monthly mean maximum plus minimum. The difference in the result obtained, however, is rarely as much as one degree. In 1933, for example, the difference was over one-half of one degree for one month only.

That marked differences in precipitation can occur between two such closely adjacent localities in Connecticut is indicated in the tables, particularly in the data for 1934. During six separate months of the year the Mount Carmel Station showed more than an inch of rainfall in excess of that shown in New Haven, and in two of those months the excess was over two inches. Nor is this the whole story. Local summer rains are of common occurrence in the State and may be limited in extent. For example, from the eleventh to the fourteenth of June, 1934, rainfall occurred every day at Mount Carmel and the total for the four days was 1.81 inches. At New Haven it rained on the twelfth only and .59 inches were recorded. On July 8 of the same year, .52 inches of rain fell at Mount Carmel and .10 at New Haven. On July 28, .53 inches fell at Mount Carmel and 1.03 inches at New Haven.

The total wind movement at New Haven during the three years was considerably in excess of that at Mount Carmel. The data are: for 1932, 82,847 and 64,381 miles; for 1933, 82,495 and 63,153 miles; for 1934, 79,181 and 56,047 miles, respectively. This same relative difference holds throughout the months of the year.

There follows an account of the work of the several departments, dealing especially with items of particular interest and importance. A complete list of projects will be found on page 95.

Progress of the Station's Work

ANALYTICAL CHEMISTRY

Fertilizers Analyzed

One of the important services carried on by the department of Analytical Chemistry is the registration and analysis of fertilizers sold in Connecticut. The results of the inspection are given in Bulletin 365.

The past year, 1934, marks the smallest number of samples analyzed in the ten year period beginning in 1925. During this decade the peak of registrations was reached in 1930 with 439 brands. In the same year the largest number of samples were examined in the laboratory. This is the natural reflection of conditions throughout the country. According to data compiled by the National Fertilizer Association, the peak of fertilizer consumption in the United States, a little more than 8,000,000 tons, was reached in 1930. It fell off nearly one-half in the next two years but is now on the increase.

In the past ten years, there has been a marked improvement in the proportion of samples meeting their guaranties. In 1923, only 55 per cent of those examined met guaranties in all respects. Since 1928, the proportion has not been less than 70 per cent; in 1930 it was 80 per cent; and during the past year it was 79 per cent.

Since 1929, commercial deficiencies exceeding \$1.00 per ton were found in not more than 3 per cent of the samples of mixed fertilizers examined. During the past year this percentage was 2.3.

Two instances of ground bone adulterated with rock phosphate were found.

More Feed Samples Meet Guaranties

During 1933, 1299 samples of feeds were analyzed and the annual report of inspection was issued as Bulletin 362 in April, 1934. Of these, 815 were official samples of commercial feeding stuffs collected by the Station Agent; 385 were samples of experimental plant materials examined for the Storrs Station; and the remainder were miscellaneous feeds examined for purchasers. The proportion of samples meeting guaranties has increased noticeably in the last three years. The proportion meeting individual guaranties has been high since the present statute concerning feeding stuffs control was enacted in 1925, but especially so since 1930.

A summary of deficiencies found on analysis of official samples, given below, is self-explanatory.

Year	No. of official samples	Samples meeting guaranties %	Individual guaranties met %
1925	488	83	94
1926-27	670	83	94
1928	716	78	92
1929	646	82	94
1930	678	81	93
1931	739	88	96
1932	836	85	95
1933	815	88	96

**17% Food and
Drug Samples
Below Par**

Examination of 1,528 samples of foods and drugs in 1933 revealed that 17 per cent fell below standard, were adulterated or otherwise illegal. This work is summarized in Bulletin 363 and was done for purposes of food and drug control in the State. Malt beverages, vegetable food oils, ice cream and spray residue on fruits and vegetables were the items of foods given most attention. No significant excesses of lead or of arsenic were found on any of the market samples examined. Drug inspection included 24 official preparations, listed in the United States Pharmacopoeia and National Formulary, of which about 250 samples were taken at drug stores throughout the State. Variations from standard quality were noted chiefly in chlorinated soda, compound solution of iodine and solution of magnesium citrate. Eight samples of whiskey labelled as medicinal grade did not meet the specifications for that article as judged by U.S.P. standards.

More than 2,000 pieces of glassware used in testing milk and cream by the Babcock test were checked and certified as to accuracy.

Members of the staff of the department of analytical chemistry have collaborated in studies of analytical methods sponsored by the Association of Official Agricultural Chemists. Dr. Fisher has investigated methods for the determination of calcium gluconate. Dr. Bailey has continued to serve on the Council for Foods of the American Medical Association.

BIOCHEMISTRY

**Chemistry of the
Tobacco Plant**

The chemical investigations of the tobacco plant carried on by the department of Biochemistry this year have included a detailed study of the rate of growth of shade-grown tobacco from the seedling to an advanced stage of maturity. Collections were made at frequent intervals throughout the season and the leaves, stems, and fruit have been analyzed chemically in considerable detail. The data obtained permit us to follow the accumulation of various organic and inorganic constituents in the plant as growth progresses; particular attention has been paid to the different forms of nitrogen and to the organic acids.

An important outcome of this work has been the study of the amide nitrogen of the tobacco plant. At least two different amides are present in tobacco, one of which is definitely known to be asparagine. There is much evidence that glutamine is also present and there are indications of still a third type of amide nitrogen.

**Protein
Chemistry**

The investigation of amides has naturally led to a detailed study of the extremely rare amide, glutamine. This substance has been found in the leaf and stem tissue of the tomato plant and is especially plentiful in beet roots. A method developed in the laboratory this year makes it possible to obtain considerable quantities of glutamine from this source in pure form. A study has also been made of the effect of liberal fertilization with ammonium sulfate upon the glutamine content of beet roots.

The studies of protein composition have included a new determination of the basic amino acids of casein and of the cystine content of the hemoglobins of the horse, sheep, and dog. The mercuric chloride complex compounds of several amino acids have been investigated with the object of developing new methods of separation of amino acids by the use of mercury reagents.

Nutrition Investigations

One of the current trends of interest in nutrition is concerned with the part played by the inorganic salts in the diet. For several years the department has carried out an elaborate series of experiments to determine the mineral requirements of the albino rat. Four of the artificial salt mixtures widely used in experimental diets employed in nutrition studies have been compared at various percentage levels in otherwise adequate rations. On the basis of rate of growth (from 60 to 200 grams body weight, as well as of percentage of bone ash) the different salt mixtures showed definite variations in efficiency, particularly at the lower levels.

Following these observations, the various salt mixtures were supplemented with pure salts. By this device it was found wherein the inferiority of the salt mixtures lay and also the reason for the superiority of the better salts. Among other things, this experiment again emphasized the significance of an adequate level of calcium in the diet of the albino rat.

With the foregoing experience available, it then became possible to formulate a new salt mixture combining the desirable features of the various mixtures now used. This extensive investigation, carried out with meticulous attention to experimental details, is another of the fundamental contributions of this laboratory to the basic nutrition of the extraordinarily useful albino rat.

Mating and Productivity of Rats

The investigation of the reproduction of the albino rat on a presumably complete diet has been continued in our laboratory for a period of four years under the supervision of Professors Arthur H. Smith and William E. Anderson of Yale University. Breeding experiments with the first four generations have been concluded.

In regard to the relationship of the interval between matings and reproduction performance, the supplementary information obtained during the past year indicates that the longer period of rest results in greater reproductive success. Whereas in the fourth generation 91 per cent of the matings were fertile in the group with three weeks of rest between periods of reproduction, only 78 per cent of the matings were fertile in the group permitted to rest but one week. Furthermore, in the former group, 80 per cent of the young were successfully weaned, whereas, in the second group, only 61 per cent of the young were weaned.

In last year's report attention was called to the high average level of body weight of young at weaning in all groups in the first three generations. A comparison at the present time of similar values in the fourth generation—based on observations of approximately 1200 animals—indicates an unmistakable trend toward lower weaning weights in all groups. These additional data emphasize the importance of carefully controlled breeding experiments extending over relatively long periods.

Further investigation of vitamin concentrates has been made. As a new feature of this work, a study has been undertaken of the rate of growth of the offspring of rats daily injected with an extract of the thymus glands of calves.

The investigations in this laboratory are supported, in part, by the Carnegie Institution of Washington.

BOTANY

Chestnut Blight Makes Less Progress

The chestnut blight (*Endothia parasitica*) seems to have made less deadly progress on seedlings in 1934. Each year for the past eight seasons, the Station Botanist with the coöperation of the Forester has planted nuts from southwestern Virginia at the Station and seedlings have been set out in selected places at Fairfield, Orange, Rainbow and Portland. A bushel of chestnuts was planted last autumn and next spring about 1,000 one-year seedlings will be ready for transplanting.

The purpose is to watch the progress of the blight; to determine, if possible, whether it is less prevalent, or whether the disease becomes less virulent, as time goes by. In the last few years more of the young trees have been killed by drought than by blight, the Botanist reports. Nevertheless the blight seems to attack them when they reach a certain size. The largest saplings are about 25 feet tall with a four-inch stem.

The past season was favorable to chestnut growth. There were many finds of burrs on sprouts from old stumps and on seedlings, and the trees increased in size because moisture conditions were better. This was especially true at Orange and Rainbow where fertilizer was applied last spring.

The chestnut tree at Lebanon, more than fifty years old and the oldest known chestnut in the State, is still alive although badly affected. The Botanist reports that it bore a number of burrs and leaves this summer and will probably survive for a year or two longer.

"X" Disease of Peach Trees

The strange trouble that appeared prominently in Connecticut peach orchards in 1932 has been receiving even more of the attention of station botanists this year. So far the exact cause has not been determined and we have called it the "X" disease.

The trouble is characterized by a premature yellowing and ripening of the foliage on all or part of the branches at midseason or later. Sections of the leaf tissue seem to be cut off from the rest of the leaf and fall out, until the foliage presents a ragged appearance. Both fruit and leaves on injured branches, with the exception of tip leaves, drop early, but the new buds on the branch develop normally, showing no symptoms of disease until the latter part of the following June. The peaches that remain on the tree seem to be normal but perhaps ripen a little early.

In the wood of injured branches there is a characteristic brown streaking, extending longitudinally through the branch, and frequently arising from a cut end of a limb, or a cut-off side branch. These symptoms were

obscured during the past summer because much of the wood of peach trees was blackened as a result of last winter's severe cold. In winter, the twig and bud growth on affected branches appear normal and it is very difficult to distinguish healthy from injured trees by external symptoms. Therefore most of the work on the peach trouble must be done in warm weather.

The Station has made investigations of the peach trouble along three lines. There were careful observations and studies in the field to find the distribution, rate of spread, extent of injury and possible control. Attempts were made to discover the nature and cause of the trouble through examination of specimens in the laboratory, budding experiments and chemical analyses. The Soils Department analyzed samples of soils from orchards scouted by the botanists, as well as from other typical orchards, to see if there was any relation between soil conditions and the trouble. Results of some of these studies are now available as recorded below:

Of the 70 peach orchards surveyed by the Botany Department in 1934, affected trees were found in 29, all of them north of Southington. The disease appeared in old and new orchards alike, and in trees planted on soil just converted to orchard use as well as in old soil. Nine orchards mapped in 1933 were checked and eight new orchards mapped. Using these maps, several growers are cutting out diseased trees this winter. In old orchards checked, it was found that the "X" disease increased from 5 to 10 per cent during the year.

In an effort to see whether the disease could be controlled by surgery, last spring unhealthy branches were cut from 41 two-year old trees and the wounds sealed with grafting wax. Of these, 44 per cent appeared sound at the end of the season, but the disease recurred in 56 per cent of the cases. Data will be taken on older trees next year.

Hundreds of specimens of diseased trees were examined in the laboratory in attempts to discover the nature and cause of the peach trouble. Cultures of several fungi were obtained from the wood and these were used to inoculate healthy peach trees to determine whether or not they would reproduce the disease. One fungus, most commonly found, produced definite streaks extending from one to six inches beyond the point of inoculation and the fungus inoculated was recovered. However, no conclusions can be reported without further research.

In 1933, budding experiments were made to see whether the trouble was caused by a virus. Buds from diseased branches were grafted into 96 healthy seedlings. Of these, 43 buds took and only three showed leaf symptoms of the peach trouble. In general, if a bud affected by a virus disease is grafted into a healthy tree, it will give the disease to the tree whether the bud grows or not. Indications are, therefore, that "X" is not a virus disease.

The Soils Department found no relation between the peach trouble and prevalent condition of soils in the peach orchards of Connecticut. The findings are given later in this report.

Although actual cause and control of the "X" disease are still a mystery, in general it seems advisable to remove affected trees from peach orchards as a precaution against spread.

**Bordeaux Doubles
Potato Crop**

The yield of potatoes may be doubled by the proper use of Bordeaux mixture, according to results obtained for four successive years at the station farm at Mount Carmel. The same experiments prove that 8-8-50 Bordeaux gives better crops than the 4-4-50 mixture. Tests this year indicate further that the amount of spray is more important than the pressure in obtaining necessary coverage of foliage.

**Controlling Late
Blight of
Tomatoes**

Bordeaux mixture and red-copper-oxide sprays have been found the most effective of materials tested in checking late blight on tomatoes. Early and thorough spraying before the blight started kept the plants free from blight for several weeks. Later applications of Bordeaux or any other materials did not check the blight once it was established. Red-copper-oxide not only was very satisfactory in control of late blight but it caused no discoloration of the tomato fruit.

Greenhouse experiments showed that the same disease, *Phytophthora infestans*, attacks potatoes as well as tomatoes. Search of the old infected parts of both hosts failed to show any sign that oöspores were developed. However, it was concluded that not only the tubers of the potato carry the mycelium but that the seeds of the tomato can also carry them, in some cases over the winter. Just what happens later with this mycelium is not known, but first infections this year and last were seen on the leaves in contact with the ground.

In 1934 the blight got an early start at Guilford, making its first appearance about the middle of August. Closely watched plots in Hamden did not show the first evidence until September. Mulching the plants or staking them off the ground failed to prevent the fungus appearing on either potatoes or tomatoes. In fact, the two plots so treated at Hamden became as quickly and as badly infected as those plots where plants lopped over the ground. Thorough spraying with Bordeaux mixture, on both plants and the ground, was very successful, and, to a less extent, frequent dusting paid.

**Studies of
Vegetable
Diseases**

The Botany Department has carried out further experiments in the control of downy mildew of cucurbits, Stewart's bacterial wilt of corn, and club-root of cabbage and cauliflower. Studies have been made of a mosaic disease of rutabagas and of an undetermined trouble of carrots.

In the greenhouse, experiments have been conducted for the control of damping-off of vegetables and other seedlings. Comparative tests have been made of various seed and soil treatments for the control of pre-emergence and post-emergence damping off, together with culture of seedlings in pure sand to which nutrient solution has been added. More seedlings survived through the sand culture method than through other treatments in soil.

**Bacterial Wilt of
Corn Less
Troublesome**

While bacterial wilt on corn (Stewart's disease) was a serious menace to the crops for the past two seasons, this year there was very little disease in the State. A member of the United States Department of Agriculture working in New England claims that the severe winter reduced infection. Either the insect carrier may have been killed off, or the bacteria may

have been destroyed in the ground. At any rate, little disease appeared on either the most susceptible (Golden Gem) or the most resistant (Golden Cross Bantam) varieties of corn. However, the Bantam showed greater resistance to smut.

Disease Survey Continues

While the *Plant Pest Handbook* covers plant troubles studied by the Station Botanist for 32 years in Connecticut, the disease survey continues for the purpose of bringing information up-to-date yearly, and reporting new diseases and injuries, or old troubles attacking new hosts. Two new troubles were listed during the past year. The most important of these is a nematode injury caused by *Aphelenchoides fragariae* on chrysanthemums, found in a nursery at Bristol. It was especially bad on the variety "Mercury". The other trouble is the less injurious fungus of sweet potato caused by *Monilochaetes infusans*.

Losses Result From Severe Cold

The severe winter of 1933-34 injured or killed a great variety of less hardy shrubs and trees, especially privet, roses, peaches and certain varieties of apple. The moist weather of early spring and fall was rather favorable to fungous troubles. The black spot of elm (*Gnomonia ulmea*), rust of apples (*Roestelia pyrata*), and late blight of tomato, and to a much less extent, of potatoes (*Phytophthora infestans*) were the most important.

Seed Testing Service

As in other years, there were direct services to farmers. Whenever growers reported vegetable crop troubles, examination and diagnosis were made and advice on control methods offered when possible. In accordance with the seed law, 156 samples of seeds were collected and analyzed and a report made to the Commissioner of Agriculture. In addition, analyses were made of 247 samples submitted by individuals, and of 91 straight seeds and 65 lawn mixtures.

Radio Spray and Weather Reports

A daily radio spray service and weather report was conducted over three radio stations from May 1 to June 15. This service, maintained in coöperation with the Connecticut State College Extension Service, gave timely information on spraying with special reference to control of apple scab, basing the recommendations on the discharge of ascospores as determined by examination of material collected in various sections of the State.

ENTOMOLOGY

Control of Oriental Fruit Moth

The search for sprays and parasites to control the Oriental fruit moth has continued actively at the Station. This is the most destructive insect pest of peach and quince in Connecticut. Sprays of derris and lead arsenate used on quinces produced a crop freer from infestation than in any previous year. In an experiment aiming to prevent moths from laying eggs on trees, one tree

was covered with tobacco cotton cloth about the first of August. Apparently the results were good.

As in previous years, parasites of the Oriental fruit moth were reared in the entomological laboratory and distributed for release in orchards. This is a cooperative program initiated and sponsored by the Connecticut Pomological Society. By a new arrangement started in 1934, the society takes responsibility for assembling the orders for *Trichogramma* parasites and collects the charge made. As peach fruit buds were all killed by the low temperatures of the winter, many growers did not think it necessary to obtain parasites. This, together with the charge, although this was very low, doubtless explains the smaller demand—10,000,000 in 1934 as against 28,300,000 in 1933.

However, the distribution of *Macrocentrus* parasites increased six-fold. A new method for rearing these, devised at the Station, made it possible to send out 24,000 in 1934 as against 4,600 in 1933. In addition 4,700 other parasites were reared and liberated in Connecticut peach orchards. These and the *Macrocentrus* are still on an experimental basis and are distributed without charge.

The Federal Bureau of Entomology and Plant Quarantine has supplied some new species of parasites from the Orient, and the staff is studying their characteristics and life habits. Some of these will be ready to send out next year. It is believed that they offer considerable promise in the control of the Oriental fruit moth in Connecticut.

Substitutes for Lead Arsenate	Experiments with substitutes for lead arsenate continue to interest both the farmer and the housewife. In recent years growers of fruits and vegetables have been confronted by a double problem: The grower must protect his crops from insect or disease injury; the consuming public is increasingly conscious of the danger in residues from poison sprays and dusts used as control measures. The Secretary of Agriculture has set limits, or tolerances, on the amount of poison permitted on foods sold interstate.
--	--

Lead arsenate is the poison most commonly employed. Both lead and arsenic are poisonous to man if taken in large enough quantities. Of the two, lead is now considered more dangerous because of its cumulative effect. In 1934, the heavy autumn rains in Connecticut washed most residue from fresh food products. Of the samples examined in the analytical laboratory, very few contained poisons above tolerance and only one of these had been offered for sale. The rest were submitted by growers anxious to know the results from certain sprays.

Experiments with substitutes at the Station followed two lines: (1) to protect foliage from arsenic burn resulting from decomposition of calcium arsenate; (2) continued experiments with cryolite, the most promising of the non-arsenicals tested last year. It was apparent that improvement in the standardization of physical properties of synthetic cryolite is greatly needed. One of the most striking characteristics is its poor adherence to the fruit. The lot used was ineffective in controlling curculio and codling moth but showed promise in control of the apple maggot.

**Attempt to
Control Onion
Thrips**

In experiments to control onion thrips, 40 pounds of finely ground, crude, chipped naphthalene were mixed with 60 pounds of hydrated lime and dusted on onion plants at the rate of about 125 pounds to the acre. The first application seemed to reduce the thrips population about 50 per cent, but later the plants were just as badly infested as the checks. The plants also suffered some injury from the treatment. These experiments will be expanded and continued.

**Corn Borer
Problem
Increases**

The European corn borer is causing heavy losses to farmers and control presents a serious problem. Search for suitable parasites and sprays is going on constantly and offers considerable hope. The "clean-up" is still our main reliance. However, to be effective it must be thorough. To obtain this without the whole-hearted coöperation of entire communities is humanly impossible.

The wet spring weather in 1934 postponed plowing of some fields until after April 10, the time limit set for clean-up of fields in the Statute. Inspectors began work in the four southern counties of the State on April 18 and continued northward. Even at that late date, time extensions had to be granted because of wet land. More careful check was made on cornfields than ever before and it was necessary to report 20 cases to the public prosecutors.

Heavy damage by the first generation corn borers in early sweet corn occurred in the truck crop areas near Hartford, New Haven, and at certain points in New London, Windham and Litchfield counties. There was an estimated loss of \$17,843 on 179 farms surveyed. The average loss per acre in sweet corn on these farms was \$48.61.

Late in the season it became evident that weed infestation has an important bearing on clean-up work, and a survey was undertaken, particularly in East Hartford. In certain upland areas, the weeds in the corn fields contained as many as 214,573 borers per acre, and on wasteland adjoining the corn fields, the borer population per acre in the weeds was 83,440. On the river meadows, corn stalks had 219,030; weeds in the fields, 106,480; and weeds adjoining the fields, 38,720 borers per acre. The weeds were mostly smart weed, pig weed and lamb's quarters. Giant ragweed on waste areas near the river contained 24,200 borers per acre.

**Mexican Bean
Beetle Studies
Completed**

The Mexican bean beetle has been a subject for active research for a number of years. The project was completed last summer and conclusions are now being prepared for publication. This insect, a lady beetle, broadly oval, pale brown in color and with spotted wing covers, is the most destructive pest of beans in Connecticut. During the past year control experiments included the effect of spacing plants, of using non-poisonous insecticides to avoid poison residue, and of comparing sprays with dusts.

Bountiful string beans were planted with 2, 4, 6, and 8 inch spacing between them in the row. Results bore out findings in previous tests. In the 2-inch spacing, there is the largest yield of pods but the largest

percentage of these is injured. The best all-round spacing was found to be 4 inches.

Insecticide experiments on Bountiful string beans showed that satisfactory results can be obtained with non-poisonous materials. Thus 25 and 50 per cent pyrethrum dusts, and .4 and .6 per cent rotenone dusts, all gave larger yields and a larger percentage of uninjured pods than magnesium arsenate spray, and an equally large percentage of marketable pods. Both pyrethrum and rotenone dusts gave good control of leafhoppers as well as of bean beetles.

Fordhook dwarf lima beans were treated with copper-calcium arsenate dust, and with Bordeaux mixture spray, in comparison with check, or untreated plots. Both treatments increased the yield about equally over that of the untreated checks but plants treated with copper-calcium arsenate dust showed the least injury from Mexican bean beetle.

	Check	Copper-Calcium Arsenate Dust	Bordeaux Mixture Spray
Yields, ounces per plant	12.1	15.1	15

More Termite Infestations Investigated

An increasing number of eastern subterranean termites, common in stumps and logs in Connecticut woodlands for many years, seem to be infesting houses of the State. In the past spring and summer entomologists examined 46 houses and found the insects in 39 of them. There were only seven similar cases reported in 1932 and thirteen in 1933.

Termites are common in tropical climates where they destroy property by hollowing out beams, woodwork and even furniture. In this part of the country the injury is not so extensive, but it may be serious and cause considerable damage and expense.

Estimated cost of repairs and termite-resistant construction to 24 of the houses examined was placed at \$6,910. The amount of money actually spent was \$3,340, since owners preferred to take care of present infestations without making other parts of their premises termite-proof.

In general, termites enter houses through direct contact between soil and wood. They may find cracks in concrete foundations and brick work, however, and sometimes build runways for themselves over a stone surface if the distance between soil and wood is not too wide. They cannot exist without wood and moisture and the workers die when exposed to the sun and air.

Of the houses examined, 15 entries were made through direct contact between wood and soil. In 11 other places, termites had found cracks in the foundation work through which they could enter wood. A complete review of findings and recommendations will be given in the report of the State Entomologist next spring.

Mothproofing Materials Tested

Moths and moth injuries constitute a perennial problem of the home maker. Experiments with mothproofing materials were continued in 1934, and the results will be published in the Entomologist's report.

**The Potato
Flea Beetle**

The flea beetle causes greater injury to potatoes than is generally appreciated. Comparative treatments on potatoes with sprays and dusts for its control this year yielded significant results. A large series of plots of Irish Cobblers was (1) sprayed with Bordeaux mixture plus lead arsenate-fish oil, (2) dusted with barium fluosilicate dust, and (3) left untreated as checks. Four applications were made. The lead arsenate-fish oil plots gave the highest yield, averaging 225 bushels an acre. Bordeaux mixture brought the second largest crop, 217 bushels, and the untreated rows, 149. Thus the highest yield was more than 50 per cent larger than the check. Moreover, lead arsenate and fish oil was the cheapest treatment.

About 25 plots of Green Mountain potatoes were included in a similar experiment comparing (1) 6-6-50 Bordeaux mixture plus 1 1/2 pounds of calcium arsenate, eight applications; (2) barium fluosilicate dust (1-3), two applications, followed by 6-6-50 Bordeaux mixture, seven applications; (3) barium fluosilicate dust (1-3), six applications; and (4) 6-6-50 Bordeaux mixture, eight applications. The highest yield, at the rate of 567 bushels per acre, was from plots treated with (2), the combination of Bordeaux and barium fluosilicate. However, this is the most expensive treatment. Straight Bordeaux (4) came second with 538 bushels; Bordeaux with calcium arsenate (1) was third with 463 bushels; barium fluosilicate dust (3) the cheapest application, yielded only 272 bushels; and untreated plots, 261.

**Relief Funds
Push Mosquito
Elimination**

The past year has seen tremendous advances in mosquito elimination work throughout Connecticut. Over a period of 29 years, 1904 to 1933, 11,000 of the 20,000 acres of salt marsh breeding places in the State had been ditched and approved for state maintenance. The ditching of the remaining 9,000 acres was nearly completed in 1934. Small areas are still unfinished in Milford, North Haven and Old Saybrook, and a larger tract in Stratford may require another season for completion. But it is estimated that the work as a whole has been pushed forward by about 15 years.

These important extensions were made possible by coöperation among the local, state and federal offices. All initial ditching in the past was done at local expense. When an area had been properly treated, the Director of the Station accepted it for state maintenance and yearly the State appropriated a small sum for upkeep. Unfortunately communities were not mosquito-proof by virtue of cleaning up their own marshes. Too frequently pests breeding in adjacent unditched territory flew, or were blown by the wind, across the border. It became evident that all marshes had to be drained if the work was to be effective. Under the circumstances progress was slow.

The bright aspect of the program in 1934 was the availability of relief funds for mosquito projects. During January, more than 1000 men were assigned from the lists of the local unemployed and were paid by CWA or state relief funds. Boots, tools and any materials necessary were supplied by the individual towns. The number of workers had been reduced to about 500 in October.

Altogether 1,476,756 feet (280 miles) of ditches have been installed and 259,825 feet (49 miles) of stream banks cleaned. Many improvements in the way of new tide gates, dikes and outlets have been made. The total federal funds expended on wages for the unemployed until October 1, 1934, amounted to \$287,299.22, and state and town contributions, \$34,729.06.

A larger maintenance appropriation by the General Assembly will be necessary before the newly ditched area can be accepted for state maintenance.

"Fresh water" mosquito projects have been carried on in the five inland towns of Ansonia, Derby, Manchester, New Canaan and Southington, and also in several shore towns.

**Investigations of
Oil Sprays**

Work with oil sprays consisted of tests of tar distillates and cresylic acid combined with commercial spray oils. However, absence of rosy aphids this year prevented collection of good comparative data. Tests were also made with tank-mixed white oils for control of pear psylla and European red mite. For the European red mite a very satisfactory summer oil preparation was made using 1 gallon industrial white oil (80 viscosity) and 1 pound powdered skim milk in each 100 gallons of spray. The combination was quite effective on the red mite. Unfortunately, it killed some other insects that prey on the red mite itself.

**Control of
White Apple
Leafhopper**

White apple leafhoppers appeared in some numbers towards the latter part of the season and a series of sprays were tried, using nicotine sulfate with and without soap, and two pyrethrum preparations. Anabasine sulfate gave the best results this year. The use of soap apparently did not increase the kill beyond that obtained with the water solution.

**Inspection
of Nurseries
and Orchards**

As usual, members of the department of entomology have inspected all of the commercial nurseries in the State and more than 20 orchards. Again there is an increase in the number of nurseries, the total reaching 376. Last year 365 were listed.

All nurseries that had important pests to be eradicated as found by the first inspection were re-inspected before certificates were issued. A further check, late in the autumn, was made on those growing pine trees to see that the European pine shoot moth had been eliminated. The presence of this insect cannot always be detected if inspection is done in July and August. On the whole it was less prevalent than in 1933, because of the severe winter and the clipping off and burning of infested shoots the previous year.

**Gipsy Moth
Control**

The Station coöperates with the United States Department of Agriculture in combatting the gipsy moth. In general, federal crews work west of the Connecticut River, and state crews to the east. The state forces had made good progress when, beginning November 15, they were asked to supervise 340 CWA men in a state project for the protection of forests under the State Forester.

After the CWA men had been trained, they were organized into scouting crews, each with one of the experienced scouts as foreman. Although the territory assigned had not been covered, all CWA work was discontinued on February 15 and the regular scouts resumed their usual scouting duties. The CWA crews made possible the discovery of rather large infestations, mostly in forest areas, in 21 towns. Some of these infestations were later cleaned up and sprayed by station crews. Altogether, east of the Connecticut River, 11 towns were completely scouted, 28 were partially scouted, and in 20 no work was done.

State and federal forces together worked in 107 Connecticut towns, found 453 infestations, creosoted 434,410 egg-clusters, sprayed 47 separate infestations, besides 2,371 acres of woodland, and 2,861 street trees. Scouts covered 4,252 miles of roadside and 435,581 acres of woodland.

Valuable assistance was rendered by men in the CCC camps. Here the work was well organized under the direction of the camp superintendents. They scouted an additional 13,828 acres of woodland.

Inspection of Apiaries

Of the 1,420 apiaries inspected by entomologists, only 27 were infested with American foul brood and two with European foul brood in the past year. There were

7,128 colonies included in these hives.

Sprays and Dusts Compared

The object of the spraying and dusting experiments, carried on coöperatively by members of the departments of botany and entomology, is to find the most economical materials that will give best control of pests and

also overcome spray injury. This year the following combinations were used on trees in the west apple orchard at the station farm at Mount Carmel: (1) a combination of lead arsenate, lime and fish oil, with no fungicide; (2) dry lime-sulfur and lead arsenate; (3) flotation sulfur and lead arsenate; (4) magnetic sulfur and lead arsenate. Several trees were left untreated as checks. Based on previous experience in these orchards, the spraying schedules were arranged to give the maximum control with the minimum number of sprays. As in previous years, lead arsenate lime-fish oil gave good control of fungi on all varieties of apples except scab on McIntosh. The tabulated results for the various treatments follow:

Average On All Varieties

	Check	Lead arsenate, lime and fish oil	Dry lime-sulfur, lead arsenate	Flotation sulfur, lead arsenate	Magnetic sulfur, lead arsenate
Good	1.43	65.2	71.69	68.86	69.79
Curculio	74.99	31.4	25.1	5.75	26.36
Codling Moth	15.23	.13	.1461
Other chewing insects	40.04	1.91	2.70	1.60	.78
Scab	11.37	1.37	.71	.47	3.67
Rust	24.43
Blotch	38.11	.68

The Insect Survey

Collections and observations of Connecticut insects went on as usual and reports were sent each month to the Insect Pest Survey Bulletin. Many specimens are

brought or sent to the Station annually for identification. Members of the department are on continual watch for specimens not included in the

extensive station collection and for insects appearing in Connecticut for the first time. They classify and mount these, keeping records of the date and place of appearance, and the hosts. They also find out from records of other states and countries what harm the insect is capable of doing and possible methods of control. Last year the following new insects appeared in the State for the first time: The beech scale in Hartford; the Dermestid beetle in Bridgeport; and the holly leaf miner in Hartford and Newington.

FORESTRY

Blister Rust Control Extended

Four projects employing an average of 370 men throughout the year made up the enlarged program for control of white pine blister rust in Connecticut in 1934. This work was directed by the Forestry Department in coöperation with the United States Department of Agriculture and was financed chiefly by the Federal Government through the CCC, CWA, ERA, and NRA.

Blister rust has been a deadly foe of white pines for many years. Thanks to eradication measures, the prospects for control in this area are now good. It is a fungous disease depending on currant or gooseberry bushes, *Ribes*, as alternate hosts to propagate the spores. It does not spread from pine to pine, but from pine to *Ribes*, and thence to healthy pine. Blister rust has threatened a valuable state resource.

This year workers were engaged in the following ways:

They removed more than 860,000 wild *Ribes* from 73,000 acres of land. These bushes grew within infecting distance, 900 feet, of white pine stands. As a result, trees in 26 towns were protected from the rust.

They made a survey of 66 towns locating 5,500 patches of cultivated *Ribes* within the danger zone. All black currant bushes, outlawed in this and some other states, were removed, and nearly 20,000 other varieties were destroyed.

Two new nursery sanitation zones were established and nine existing zones re-checked, with the removal of 2,150 wild, and 51 cultivated, *Ribes* from 3,685 acres.

Maps showing pine stands were made in 15 towns in the northern part of the State. As a result, the efficiency of future control work in these towns should be greatly increased.

The White Pine Weevil

Another enemy of the white pine is the weevil that attacks leaders and causes distortion of the trunk, thus spoiling its timber value. Local control undertaken last year was continued from the middle of June to the early part of August by crews from nine CCC camps. More than 5,000 acres in 21 towns were covered and approximately 128,000 infested leaders removed. A comparison with last season's work shows the infestation to have decreased throughout the 10 areas covered. In one section of 150 acres, the infestation was almost 50 per cent less than the previous season. It is anticipated that work may be continued for another year,

and that better information will be secured regarding the possibility of controlling this forest pest.

**Measuring
Pine Weevil
Damage**

A second project in connection with the white pine weevil was carried on by the department of entomology, acting in coöperation with the United States Bureau of Entomology and Plant Quarantine. In this, measurements of the degree of trunk distortion in weeviled white pines of various ages and rates of growth are being made. The purpose is to develop a method of predicting the amount of recovery that will occur in weeviled trees before the crop is cut.

**Pine Shoot Moth
Infestation
Reduced**

Control measures and the severe cold of last winter seem to have reduced the European pine shoot moth to the lowest point in three years. The larva infests the tips of red, Scotch and some other pines, retarding growth. Repeated attacks may kill the trees. Between September, 1933, and July, 1934, crews from the CCC camps and CWA workers carried out control measures directed by the departments of forestry and entomology. Nearly 800 plantations comprising 11,000 acres in 125 towns were covered, and millions of infested shoots were cut off and burned. Spring measures were most effective since they followed the killing cold of the 1934 winter. It is estimated that between 80 and 90 per cent of the larvae in the shoots perished. Plans for the coming season include thorough scouting during the winter with men furnished by the ERA, followed by a vigorous program during the spring of 1935 with CCC crews. The entomologists are making a survey of all red pine plantations, and further biological data, particularly regarding parasites and predators, must be gathered next year.

**Durability of
Treated Posts
Tested**

Since the disappearance of the chestnut from the hill-sides of Connecticut, farmers have been hard pressed to find durable wood for posts. Experiments conducted by the Forestry Department have shown that other native woods properly treated with creosote may equal or even surpass the chestnut in durability. Tests were made of the service life of posts of red pine, pitch pine, Scotch pine and red maple treated by the open tank process described in previous reports. These were placed in the ground with suitable check posts under the following conditions:

Seven posts of each species—four treated and three checks—were set in a pasture fence at Middlebury in comparatively heavy soil.

Eleven posts of each species—six treated and five checks—were set in a tobacco shade tent in light, sandy soil.

All posts in the pasture fence were in good condition after two years, with the exception of one red maple check and one Scotch pine check. These showed signs of rot just above ground. In the shade tent, all the check (untreated) posts had become entirely unserviceable and had to be replaced. Without exception the treated posts remained good. It appears, therefore, that conditions under the tobacco tent favor decay but that treated posts will withstand such conditions.

**Treatment of
Tobacco
Tent Poles**

Another experiment testing the life of treated poles has been carried on for five years at the Tobacco Substation. Results of creosoting white pine, pitch pine, gray birch, red maple and popple show that: (1) Untreated poles set in the ground rot and become unserviceable in less than five years; (2) Creosote applied to the butts with a brush extends the life of the poles a short time only and cannot be recommended; (3) Treating the butts in hot creosote for three hours (open tank method) renders all species still serviceable at the end of five years; (4) The tops of the poles may be preserved by merely brushing with, or dipping in, creosote; (5) With proper treatment any of the species mentioned can be satisfactorily used for tent poles. From the standpoint of abundance, price, adaptability to treatment and physical properties, the most suitable species are red maple and pitch pine.

**Post-Treating
Plant
Established**

A commercial post-treating plant has been established by the State Forester on the Meshomasic Forest in Portland. Experimental tanks for use by the Experiment Station have been set up in connection with this plant. Although the commercial plant has been in operation for several months, there has as yet been little opportunity to use the outfit. Much experimental work will be carried on in the coming year. Therefore it seems best to delay the publication of a fuller report until these data are secured.

PLANT BREEDING

**Seed of New
Early Sweet
Corn Ready**

Seed of a number of outstanding strains of corn developed by the Genetics Department has been increased and is ready for distribution through seed companies this year. Two of these are new top crosses that have proved remarkably productive of large, well-filled ears as early in the season as the first commercial strains. Spancross C2 has been outstanding in these respects and is also highly resistant to bacterial wilt. This variety is the result of a top cross of Spanish Gold and Connecticut 2 (a Whipple inbred). A similar cross of Golden Early Market and Connecticut 6 inbred, called Marcross C6, matures a few days later but produces slightly larger ears. Spancross P39 (Spanish Gold by Purdue 39) yields a large number of medium sized, well-filled ears of excellent quality. Seed for each of these is available.

**Midseason
Sweet Corn
Tested**

As a result of several years of testing, Whipcress C6.2 appears to be most satisfactory for midseason corn. This hybrid is a first generation cross of two Whipple inbreds. The plants are uniformly productive, yielding about 10,000 marketable ears to the acre, and highly resistant to bacterial wilt. Practically the whole crop may be harvested in one day.

Adaptability and ease of producing seed recommend Whipcress P39. This is a top cross of the Whipple variety as seed parent with Purdue 39 inbred as pollen parent. It is little, if any, inferior to Golden Cross Bantam,

has a larger ear and matures three to five days earlier. It has been outstandingly productive of attractive ears of good quality; but the ears are not so uniform as in Whipcross C6.2 and are later in ripening.

Breeding Experiments Bearing on Evolution

Results of experiments crossing paired lines of inbred corn may have an important bearing on evolutionary theory. After 17 to 23 generations of self-fertilization, three inbred lines of corn were divided into two lines each and these continued by self-fertilization for five to ten generations further. Crosses between these paired lines have been made to show whether or not transmissible variations have occurred. If changes in the direction of better growth are handed down from generation to generation, the work would contribute evidence on evolution and to the analysis of growth factors.

Southern Corn High Producer Here

For the fifth successive year 1934 data confirmed previous trials showing that the late southern ensilage varieties of corn produce about 30 per cent more dry matter than the northern varieties usually grown here. Planted on the station farm at Mount Carmel, this corn grows to a height of 13 feet and green material weighed in the field amounts to about twice as much per acre as that from native ensilage. Probably this remarkable growth is not general for the whole state but only for a limited area along the coast and possibly up the Connecticut River within the upper Austral biologic zone. Most of the corn belt in the middle western states lies in this same zone. The rest of Connecticut is in a different zone, called Transitional, and in this area the southern varieties of corn would probably not give as large increases in yield as earlier varieties.

New Strain of Squash Developed

Investigators working with watermelons, field pumpkins, and cantaloupe have demonstrated that high and low yielding strains of all these cucurbits can be isolated by inbreeding and selection. These plants cross pollinate so readily when grown in the field, that in this section inbred lines can be obtained only by hand pollinating and covering the blossoms.

Our efforts with summer straightneck squash have resulted in the isolation of an early, productive, uniform strain of this variety which appears to be well adapted to New England conditions. Seed of this strain is being increased and will be available for distribution in 1936.

Better Pepper Hybrids Developed

After a number of years of work in greenhouse and field, several new hybrid peppers of the most promising quality have been developed. These are the result of crossing California Wonder with a natural hybrid from sweet Spanish pepper and Harris Early Giant. The plants are sturdy and produce prolifically and earlier than standard varieties. Fruits are of good size, blunt nosed and green, with smooth, fleshy walls. Seven tentative selections from 60 fourth generation hybrids have been made. They are to be increased and the best strain will be introduced as a new variety. In addition to the seven selections, 80 individual plants were selected for further comparison with commercial varieties in 1935.

New Straw-berries Prove Superior Approximately 400 varieties of strawberries, resulting from crosses made in 1931 and selected from 9,000 seedlings in 1932 and 1933, were fruited at Mount Carmel in 1934. Standard varieties were grown for comparison. Some of the new varieties proved superior in yield to all commercial varieties tested. About 50 of the most promising have been selected for further trial. They will be fruited at Mount Carmel in 1935. From these it is hoped to obtain a few (possibly 10 or 12) outstanding selections that will be propagated and tried out by the farmers of Connecticut and adjacent states.

Lima Beans A pure line selection of the Fordhook bush lima bean has been tested for three years and shown to be more productive than commercial stocks of the same variety. There is a question whether this is due to an inherited difference or to the manner of growing seed. Commercial seed of lima beans is frequently injured in threshing. Being shelled by hand, the locally grown beans may have an advantage. This point is being tested.

New York Lettuce Pure line selections of New York lettuce have failed to show consistent improvement over the original strains. The variation within the pure line far exceeds the variation among the different lines. This fact, together with the difficulty of obtaining seed in Connecticut, makes the improvement of this important vegetable crop exceedingly difficult. Crosses with other varieties have so far failed to bring out desirable heading qualities.

The variation in per cent of marketable heads of different commercial strains of lettuce during different seasons due to soil and weather conditions tends to obscure any inherent differences in the seed itself. Recommendations based on limited trials have doubtful value.

Crossed Tomato Plants Individual plants have been selected for four generations from crosses involving Alacrity x Bonny Best x several of the standard commercial varieties. Fourteen of these selections were compared with several commercial strains this year. The best selection gave a total yield of 52,969 pounds per acre. In comparison, the next highest yielding variety, Penn State Earliana, gave 50,268 pounds. The average weight of individual fruits on the selection was 4.2 ounces, a weight highly desirable on the market. Some of the commercial varieties tested weighed more, and some less, than this. The work will be continued.

SOILS

Soil and Land Cover Inventory The department of Soils has worked out a practical plan for mapping soils and land use throughout Connecticut by the use of the air photographs now available. During the spring of 1934, a complete airplane survey of the State was made as a project of the State Planning Board. Several years ago, the Station used an air map of Middletown as a base

in mapping the soils and land cover of that area. The experiment proved definitely that much more accurate field work may be done, in materially less time, if the area has been photographed first. Since then, several other states have experimented with air photographs in soil mapping and a considerable body of experience is now available.

With a view to undertaking a complete soil and land use survey of the entire state in the near future, an area of 30 square miles in the town of Glastonbury was selected for study of mapping technique, using the new aerophotographic mosaic (scale 1 inch = 2640 feet or 1/2 mile). As a result, a rapid practicable method of accurate mapping of soils and land use has been devised.

Increased interest in land use planning makes such a survey necessary. Planning includes not only more intelligent adjustment of agriculture to soil and topographic conditions, but the development of recreational areas, state forests, game preserves and part-time farms. A survey cannot be made with available research funds, but could be undertaken at minimum expense to the State through coöperation with federal relief agencies and the soil survey division of the United States Bureau of Chemistry and Soils. It is hoped that the work will start in 1935.

Soil Testing Helpful to Farmers

The "Universal" system of soil testing developed by this Station is proving helpful to hundreds of persons who submit samples in the course of each year. The system is constantly being standardized with the results of pot tests, fertilizer experiments and with field observations on large numbers of samples tested. The interpretations of the tests are now much improved and the Station offers a valuable service to citizens of Connecticut. Our system is now being used by numerous experiment stations, extension services and commercial agencies.

Peach Soils of Connecticut

The appearance of the baffling "X" disease of peaches two years ago led to a study of peach soils in healthy as well as in affected orchards. While no relationship between the nutritional characteristics of the soils and the occurrence of the new peach trouble has been evident, it is believed that more attention should be given to liming and fertilization in the orchards of the State, especially if satisfactory green manure and cover crops are to be grown. It is true that apparently good orchards are growing on strongly acid soils of low mineral fertility, yet it is reasonable to suppose that these trees would be better off in the future if a more favorable condition of the soil were maintained.

A series of 207 soil samples from 56 commercial peach orchards showed wide variations with respect to soil acidity and the availability of the mineral elements related to plant nutrition. The peach trouble occurred on 64 of these samples. A prevalence of strongly acid conditions and generally low availability of the plant food elements were observed.

Forty-nine per cent of the samples were more acid than 5.2 pH, and only 27 per cent showed reactions favorable for the growth of leguminous crops (5.6 pH or higher). This generally acid condition was also evidenced by unusually high active aluminum concentration, a factor which is correlated with the injurious effects of soil acidity. Phosphorus tests on 60

per cent of the samples were below levels usually accepted as normal for good agricultural soils. Potassium tests were low on 53 per cent of the samples.

More detailed chemical studies and pot experiments on 13 soils from representative orchards gave similar conclusions. Only three of these soils failed to give striking responses to phosphorus, potassium or lime, when tobacco, tomatoes and sweet corn were grown as indicator crops. All of the soils showed consistent response to nitrogen, but to a greater degree on the unlimed treatments. One soil was sufficiently acid to produce injury from excessive manganese concentration, while four soils showed striking symptoms of magnesium deficiency, when tomatoes were grown.

**Market Garden
Fertilizer
Trials**

The vegetable fertilizer experiment on the Windsor field at the Tobacco Substation has now been under way for five years. During 1934 growing conditions were especially favorable for the early crops. An early planting of lettuce showed best results from highest rate of nitrogen application (135 pounds per acre). On the later planting, maturing heads in warmer weather, a much smaller amount of nitrogen (45 pounds per acre) gave most compact heads, but the high nitrogen was distinctly detrimental to heading. As in previous years, manure alone, even in heavy application, has not been adequate for early vegetables. This is due to the slow availability of the nitrogen in manure during the spring months.

As a result of the intensive cropping that has been practiced on this field, several crops are beginning to show increased yields for the heavier rate of potash application (180 pounds K_2O per acre). During earlier years of the experiment a smaller amount of potash (120 pounds K_2O per acre) was sufficient.

**Fertilizing
Sweet Potatoes**

The Connecticut Valley is showing increased interest in sweet potatoes and this crop is now included in the fertilizer trials at Windsor. It has been found that best quality potatoes are grown on plots receiving a low rate of nitrogen (45 pounds per acre) and high potash (180 pounds per acre). The yields of plots so treated were 410 and 443 bushels per acre respectively, in 1934. The manure treatment, with half-rate fertilization, produced the highest yields, 594.5 bushels per acre, but the crop was irregular in size and shape. In the light of these results and the experience of sweet potato growers in the main commercial districts of other states with similar soils, it is believed that manure should not be used just prior to this crop, and that the fertilizer should supply not more than 50 pounds of nitrogen, and not less than 150 pounds of potash, per acre.

**Potatoes on
Old Tobacco
Land**

For the second year potatoes have been grown at Windsor on land previously used for tobacco. The object is to find the most satisfactory rate of application of various fertilizer constituents on such soil. Due to dry weather, the average yield for the standard treatment in 1934 was only 273 bushels per acre, compared with 376 bushels in 1933. Therefore the results this year are not very significant.

However, the residual effect of former heavy fertilization for tobacco was still in evidence. The average yield without phosphoric acid (271 bushels) was only two bushels less than the standard (273 bushels) with 120 pounds of P_2O_5 , while the no potash treatment gave only 254.5 bushels per acre. With 60 pounds of nitrogen per acre, the yield was 255.1 bushels, indicating that during a year of low yields the full rate of application of nitrogen (100 pounds) is not entirely utilized. The plot receiving no fertilizer of any kind during both years yielded 160 bushels per acre in 1934 as against 279.4 bushels in 1933. Additions of magnesia in 1933 gave some increases. However, unfavorable weather in 1934 made the yields so low that the soil, although relatively low in available magnesium, furnished sufficient of this element for the crop.

**Acid or Alkaline
Effects of
Fertilizers**

During recent years there has been a lively interest in the acid or alkaline tendencies of fertilizers, particularly those containing nitrogen. Much of the nitrogen in mixed fertilizers is now supplied as sulfate of ammonia or ammoniated superphosphates, materials which are known to be acid in their effects upon soil reaction. The lysimeters at Windsor have now yielded excellent data in regard to the magnitude of these effects. As measured by the amounts of acid constituents in the uncropped lysimeters, a complete fertilizer supplying 1000 pounds of nitrogen during a five year period has caused the following net losses or gains in soil bases.

The results are stated in terms of pounds of limestone ($CaCO_3$) equivalent:

Source of Nitrogen (1000 lbs. of N)	Net Change in Base Status of Soil in Lbs. $CaCO_3$
Nitrate of Soda	1095 gain
Sulfate of ammonia	3470 loss
Urea	1330 loss
Cottonseed meal	940 loss
No nitrogen	510 gain

One of the above soils, under cropped conditions (tobacco) and similarly fertilized, showed the following results, computed on the basis of difference between bases added and removed, one-half of the nitrogen in the crop being assumed as basic:

Source of Nitrogen	Net Change in Base Status of Soil in Lbs. $CaCO_3$
Nitrate of Soda	1533 gain
Sulfate of ammonia	3979 loss
Urea	721 loss
Cottonseed meal	1025 loss
No nitrogen	771 gain

One of the soils studied for five years is Wethersfield loam. At the beginning of the experiment this soil was well supplied with bases and only moderately acid. The effect of the acid type fertilizers on this soil could be calculated from the fertilizer formula. The other soils, strongly acid when the treatments were begun, were depleted in active bases to a significantly less extent than would be expected from the analyses of the drainage waters. On these soils much of the acidity produced by the

fertilizers was washed from the soil in combination with aluminum, manganese, and bases dissolved from the soil minerals that are not active under normal conditions of reaction of the soil solution.

Lime and Acid-Reacting Fertilizer A new experiment was started in the spring of 1934 to provide data on the amount of lime required to adjust the acidity of acid-reacting fertilizers. Two soils, originally identical, were selected for tests. One was only slightly acid as a result of liming twice during the preceding five-year period. The other was strongly acid. Each was treated with sulfate of ammonia, urea, and cottonseed meal, both without lime and with amounts of lime equivalent to the theoretical acidity of the fertilizer. Nitrogen as nitrate of soda, and a "no-nitrogen" treatment also were included on each soil.

Nitrogen Recovery With a total of five yearly 200-pound applications of nitrogen, the Windsor lysimeters accounted for nitrogen in crop removal and leaching as follows:

Source of Nitrogen (1000 lbs. of N)	Nitrogen in leaching	Nitrogen in crop*	Total Nitrogen Recovered
Nitrate of soda	763.8	373.4	1137.2
Sulfate of ammonia	680.3	243.4	923.7
Urea	498.0	322.5	820.5
Cyanamid	555.6	320.1	875.7
Cottonseed meal	432.9	285.2	718.1
Dried blood	554.3	315.6	869.9
Stable manure	370.4	175.8	546.2
No nitrogen	213.2	83.5	296.7

*Four crops only were removed, the 1929 crop being destroyed by hail.

Since serious nitrogen losses by leaching did not occur during any growing season, the nitrogen taken up by the crop was in definite relation to the amount available in the soil, except for sulfate of ammonia and cyanamid. Sulfate of ammonia produced poor crops as a result of excessive soil acidity, and cyanamid proved somewhat harmful to growth at this heavy rate of application, even though applied a month in advance of planting time.

Conservation of Fertilizer by Cover Crops The growth of cover crops has continued to show significant results in conserving residual fertilizer nitrogen against leaching losses. The average saving due to the oats cover crop grown in the fall of 1933 was 53 pounds.

In 1933-34, tobacco was followed by a fall-seeded oats cover crop. When 200 pounds of nitrogen were applied in the fertilizer, 102 pounds were removed by the crop, and only 9 pounds were lost from the soil by leaching in a 30-inch depth lysimeter. Thus a saving of 89 pounds resulted from the use of this cover crop.

Lysimeter Data from Forest Soils The lysimeters installed in a red pine plantation in the spring of 1933 are yielding some interesting data. When the soil is kept bare at all times it absorbs and passes into the receiving pots only 30 to 50 per cent of the total rainfall recorded in the open; but when the soil is covered with natural litter, between 70 and 100 per cent is recovered.

At the end of the first year rainwater that passed only through the litter contained about 31 pounds of nitrogen per acre, of which 52 per cent was in the form of organic nitrogen, 26.8 per cent as nitrates and 21.2 per cent as ammonia. Leachate from the bare soil contained 36.6 pounds per acre of which 91 per cent was in nitrate form and only 3.3 per cent was ammonia. Where the rainfall passed through both litter and soil, it contained 63.1 pounds of nitrogen, with 87 per cent as nitrates and 7 per cent as ammonia. The late summer and fall period yielded more nitrogen than either the winter or spring and early summer periods.

Recently a set of "pan type" lysimeters was installed for the purpose of studying the leachate obtained where root competition is not disturbed by the installation. Due to utilization by the tree roots, a much smaller proportion of the rainfall and considerably less nitrogen were recovered than in the case of the original cylinder type of lysimeter in which no roots are present.

Moisture in Forest Soils

Studies of soil moisture, in relation to the moisture equivalent on forest soils sampled during the driest part of the past summer, reveal variations in relative wetness in the top six inches of soil from less than 50 in the pine plantation at Rainbow, to more than 200 in a portion of the Meshomasic Forest. The average of 86 samples from 24 mixed hardwood locations was 109.1. In general, the more favorable sites as indicated by vigor of growth and stand composition had, at the time of sampling, a relative wetness of 125 or better. The lowest value found, 25.6, was obtained in a red pine transplant bed on a coarse sand at Rainbow. For comparison, the relative wetness of a garden soil on which lettuce wilted during the hottest part of the day was 137.

Effect of Removing Litter

Removal of the litter annually by raking or burning, with and without addition of lime, has not had any consistent effect upon either height or diameter growth of a 32-year-old red and white pine plantation on the Merrimac loamy sand soil at Rainbow during the first five-year period. An extra plot which received additional litter to three times its normal amount has shown some increased growth. This thick duff maintains a higher moisture content than does duff of normal thickness, but it has had no effect on the moisture content of the underlying mineral soil.

TOBACCO

Amount of Phosphorus on New Land

A five-year field test on land not previously cropped to tobacco indicates that special carriers of phosphorus, such as precipitated bone and superphosphate, are not necessary on such land. The phosphorus supplied by cottonseed meal and other organic sources of nitrogen is sufficient for the needs of the crop. High phosphorus applications have impaired, rather than improved, quality.

**Placing Fertilizer
in the Row**

Can the amount of fertilizer be reduced by applying it in the row instead of broadcasting as is usual? The results of the first year of an experiment on Shade Tobacco were not very promising. A reduction of one-quarter below the standard broadcast application gave poorer quality and yields. Even so small a reduction as one-eighth impaired the quality although the yield was satisfactory.

**Is Starter
Necessary?**

For two years Broadleaf tobacco has failed to show any advantage from so-called starter in fertilizer mixtures. Most tobacco fertilizers have a small portion of their nitrogen in the form of nitrates. The purpose is to furnish immediately available nitrogen to the young plants to insure a quicker start and more continuous growth. Results on Broadleaf field plots with and without starter have failed to show any advantage of the starter either in quicker growth or in better yield or quality.

**Quantity of
Fertilizer Nitrogen**

To apply too much nitrogen is as bad as to apply too little. A three-year field test on Havana Seed shows that plots treated with 200 pounds to the acre produced the best tobacco. Results from the use of different quantities follow:

(1) Cured leaves from the 100-pound plots are yellow, dead, chaffy and all but worthless.

(2) On the 150-pound plots these symptoms are less evident but it is clear that the quantity of nitrogen is not sufficient on this light type of soil.

(3) The 200-pound plots produced the best tobacco, with the 250-pound plots a close second.

(4) On the 300-pound plots the yield was heavier but the tobacco was coarse and showed more prominent veins, indicating excessive nitrogen.

**Effects of Different
Nitrogen Carriers**

In 1926, the Substation began experiments on four plots of Havana Seed to learn the specific effects, when used singly, of different fertilizer materials commonly used as carriers of nitrogen. The tests now include plots of both Havana Seed and Broadleaf in two other fields, added from time to time during the past eight years. Specific differences, believed by many growers to be produced by the use of certain organic carriers, either have not been evident in these tests or, if present, have not been pronounced. Cottonseed meal, castor pomace, fish meal, linseed meal and corn gluten meal, each used as the single source of nitrogen, have produced tobacco which could not be distinguished in quality or quantity one from the other.

**New Strains of
Havana Seed**

For several years a number of promising rootrot resistant strains of Havana Seed tobacco have been grown, compared and tested at the station farm at Windsor. This is in cooperation with the United States Department of Agriculture. Most of these strains proved to yield more than the standard types, are rootrot resistant and some have good quality. Similar reports came from farmers growing some of the strains on a larger scale. While they give much promise, their commercial value must wait the verdict of the cigar manufacturer.

**Station Breeds
New Cuban Shade**

By breeding and selection through a period of six years, the Station has developed some new strains of Cuban Shade tobacco. In some respects these are superior to the type usually grown, particularly in leaf shape and yield. In 1934, for the first time, a small quantity of seed was supplied to growers for larger commercial trials. Reports on these have been favorable but it is too early to learn what the cigar manufacturer's reaction will be.

**Disease Survey
of Tobacco
in 1934**

During the growing season of 1934, a large number of seed beds and fields were inspected for tobacco diseases. The season in general was characterized by less disease than any recent year. In the seed beds, only one case of wildfire and one of serious black rootrot were found. Pythium damping-off was almost absent but a number of cases of the later Pythium rootrot were observed and caused severe damage in a few beds. Serious losses from fertilizer injury were also incurred in several seed beds.

Wildfire in the field was found on one farm only. Black rootrot was not serious anywhere. Brown rootrot was confined to a few fields where tobacco had followed grass or corn. Mosaic was very much less prevalent than usual, causing some damage in a very few Broadleaf fields. Leaf spots of various types became general late in the season. Pole sweat in the sheds, aggravated by excessive dampness during the curing season, was the one serious disease of the year and ruined thousands of dollars worth of otherwise good leaf.

Investigations of leaf spots that developed late in the season showed that there were at least six different types. Two of them, wildfire and angular leaf spot, are caused by bacteria. The first appeared on one farm, while the second was more general but nowhere serious.

There were also two virus spots. The rust or fleck spots accompanying mosaic were found wherever mosaic occurred. The second virus spot, ring-spot, appeared commonly but has attracted little attention in this State. Three other types of spots, the John Williams broadleaf spot, brown spot and white speck appear to be due to physiological break-down of the leaf cells. All three were unusually abundant this season. A fungus, *Alternaria tenuis*, was found to be commonly associated with white speck and sometimes with the other physiological spots. However, considerable investigation leads us to believe that the fungus is not the primary cause of any of these spots.

**Insect Pests of
Tobacco**

Cutworms in unusual numbers infested seedbeds this year. It was found that they could be controlled by using a poison bran bait of the kind commonly used in the fields against this insect. The hornworm became uncommonly troublesome in one Broadleaf section, calling for measures more drastic than the usual hand picking. Control experiments compared arsenate of lead dust with barium fluosilicate. Either controls hornworms but treatments should begin while the larvae are small. The larger worms are not easily killed.

Comparison of barium fluosilicate with pyrethrum dust and rotenone dust for control of flea beetles showed that all three materials are effective.

However, barium fluosilicate protects leaves for a longer time and does not need to be applied as often as the other two.

The Library

During the year ended October 31, 1934, the Station Library has had approximately the following number of additions:

U. S. Department of Agriculture bulletins and reports.....	939
State Agricultural Experiment Station publications	1,354
Scientific and agricultural domestic and foreign journals (separates).....	2,722
Single books purchased.....	70
Total.....	5,085

The library subscribes to 85 sets of scientific journals. It receives in exchange for its own publications about 21 sets of domestic farm journals and 24 sets of foreign agricultural journals.

The total number of cloth and paper bound volumes on hand is now about 20,000. Most of the United States Department of Agriculture and State Experiment Station publications as well as journals are received in pamphlet form and are not included in the volume count until bound.

Botanical Collections

Dr. Clinton has given his large, personal collection of plants, botanical books, botanical periodicals, lantern slides, etc., to the Station. These include:

Books	175 to 200 volumes
Reprints of botanical articles, etc.	5,000
Bulletins, circulars, etc.	6,500
Lantern slides	several hundred
Botanical specimens	10,000
Letters from botanists	3,000

Projects for 1934-35

Analytical Chemistry

1. Inspection of fertilizers.
2. Inspection of feeding stuffs.
3. Inspection of foods and drugs.
4. Calibration of Babcock glassware and thermometers.
5. Analyses of insecticides and fungicides.
7. Analyses of special and miscellaneous foods.
8. Collaborative studies on analytical methods.

Biochemistry

1. Cell chemistry.
 - a. A detailed examination of the nitrogenous constituents of plant cells, in particular those of leaf tissues. The further development of methods for the determination of the different forms of nitrogen in extracts of such tissues.
 - b. An investigation of the constituents of the tobacco plant with special reference to the changes that occur during growth and curing.
 - c. An investigation of the composition of tobacco seed.
2. Protein chemistry.
 - a. The methods for the determination of the basic amino acids yielded by proteins with the object of increasing their accuracy and convenience.
 - b. The methods for the separation of other amino acids yielded by proteins.
 - c. The properties of certain of the amino acids and their derivatives.
 - d. Methods for the preparation of pure proteins on a large scale with the object of obtaining material for chemical and nutritional study.
3. Nutrition investigations.
 - a. The relation of diet to the rate of growth with special attention to certain factors that appear to determine rapid growth.
 - b. The investigation of the relation of certain constituents of the diet, in particular the numeral nutrients, to growth.
 - c. A study of reproduction in the Osborne-Mendell strain of white rats.
 - d. An investigation of the effect of extracts of the thymus gland on the rate of growth of the offspring.

Botany

2. The nature and cause of mosaic diseases of plants.
5. Plant disease survey of Connecticut.
8. Spraying and dusting experiments on apples and peaches. (See also Entomology, No. 3).
15. A study of the virulence of the chestnut blight.
20. Diseases of shade trees.
27. Investigations of elm diseases.
28. Studies on the identification of apple varieties by seed characters.
30. Investigations on the diseases of vegetable crops and their control.
31. Investigation of a new peach trouble ("X" disease).

Control and Service

12. Seed testing. (In coöperation with Commissioner of Agriculture).
25. Spray service (with the Extension Service, Storrs).

Entomology

3. Spraying and dusting experiments on apples and peaches. (See also Botany, No. 8).
9. Insect survey of Connecticut.
17. Studies on the control of the Oriental fruit moth.
28. Investigations on oil sprays.
30. A study of insects that attack the tobacco plant. (See also Tobacco Substation, No. 20).
31. Studies on the biology and control of the European pine shoot moth. (See also Forestry, No. 13).
32. The biology and control of the potato flea beetle.
34. Tests of methods to control clothes moths.
35. The biology and control of the white apple leafhopper.
36. The control of onion thrips.
37. Substitutes for lead arsenate in orchard sprays.
38. The relation of rate of growth and pruning methods to the recovery of white pine to weevil injury. (In coöperation with the U. S. Dept. Agri.)
39. The Carpenter ant as a pest of telephone poles.
40. Studies of sprays and parasites for the control of the European Corn borer. (In coöperation with the U. S. Dept. Agri.)

Control and Service

10. Inspection of orchards and nurseries.
11. Control of gipsy moth. (In coöperation with the U. S. Dept. Agri.)
12. Elimination of mosquito nuisance.
13. Inspection of apiaries.
19. Control of the European corn borer. (In coöperation with the U. S. Dept. Agri.)
24. Control of the Asiatic beetle.
25. Control of the Japanese beetle. (In coöperation with the U. S. Dept. Agri.)
27. Rearing and distributing parasites of the Oriental fruit moth. (In coöperation with the Conn. Pomological Society.)

Forestry

1. Experimental plantations on a sandy tract at Rainbow.
 - a. Comparison of many species of conifers and hardwoods, in pure stands and in combinations.
 - b. Methods of management for those species that have survived.
 - c. Studies on growth and habits of the several species.
2. Effect of thinning in white pine at Shaker Station.
3. Effect of thinning in hardwoods at Quassipaugh Lake.
6. Studies of forest plantations throughout the State.
 - a. Comparative growth of various species.
 - b. Reasons for success or failure.
 - c. Soil and other site factors necessary for success of each species.
10. An investigation of the distribution and growth of forest trees as influenced by soil conditions and other site factors.
11. Coniferous seed bed study to determine:
 - a. The value of fertilizers in seed beds.
 - b. The value of different amounts of seed.
 - c. The value of dusts and sprays in preventing damping off.
12. A study of preservative treatments of native woods used for posts.
13. Studies on the biology and control of the European pine shoot moth. (See also Entomology, No. 31).
14. Studies on extensive control of the white pine weevil.

Control and Service

5. Distribution of forest planting stock. (Under Clarke-McNary Act).
7. Control of white pine blister rust. (With U. S. Dept. Agr.).
15. Control of Dutch elm disease.

Genetics (Plant Breeding)

1. A genetic study of hereditary characters in corn involving their linkage relations and variability.
2. The effects of inbreeding and crossing upon corn.
3. Methods for the improvement of naturally cross-fertilized plants by selection in self-fertilized lines, with particular attention to field corn for grain and ensilage; alfalfa; some of the more important vegetable crops, such as sweet corn for market gardening and canning, beets, carrots, cucumbers, melons, squash; and some fruits, such as bush fruits and strawberries.
4. Methods for the improvement of naturally self-fertilized plants, with particular attention to tobacco and vegetable crops such as lettuce, lima beans and tomatoes.
5. A study of variation and the effects of selection in strains of cross-fertilized and self-fertilized vegetables.

Soils

1. A descriptive inventory of Connecticut soil types in relation to their use for crops, pasture and forest.
2. The physical and chemical characteristics of important soil types in relation to the nutritive response of tobacco and other crops when these soils are variously treated in the greenhouse.
3. Nutrient requirements of vegetable crops on important soil types used for market gardening in the State.
4. A study of the physical, chemical and biological conditions of several soil types in natural mixed hardwoods and in planted coniferous forests.
5. Lysimeter studies of the drainage losses and other changes that occur in several soils under heavy fertilization as practiced for tobacco and vegetables.

Tobacco Substation

1. Fertilizer experiments—various sources and rates of nitrogen, phosphoric acid, potash, lime and magnesia.
2. Field tests with farm and "commercial" manures.
4. Tobacco nutrition studies—the role of nitrogen, sulfur, potassium, calcium, manganese, boron and magnesium.
5. Improvement of Havana seed tobacco. (With U. S. Dept. Agr.).
6. Improvement of Broadleaf tobacco.
7. Improvement of Cuban shade tobacco.
8. The effect of various winter cover crops used on tobacco land.
11. Soil reaction in relation to tobacco.
13. Preservative treatment of shade tent poles. (See Forestry, No. 12).
17. The role of humidity and temperature in curing tobacco.
19. Diseases of tobacco.
20. A study of insects that attack the tobacco plant. (See also Entomology, No. 30).
22. Irrigation experiments.
23. Studies on the rate of growth of tobacco.

Publications

BULLETINS OF THE STATION

- REPORT ON COMMERCIAL FERTILIZERS FOR 1933. E. M. Bailey. No. 355.
STUDIES ON THE PARASITES OF THE ORIENTAL FRUIT MOTH. II. *Macrocentrus ancylovorus*. Philip Garman. No. 356.
REPORT OF THE DIRECTOR FOR THE YEAR ENDING OCTOBER 31, 1933. No. 357.
PLANT PEST HANDBOOK FOR CONNECTICUT. II. Diseases and Injuries. G. P. Clinton. No. 358.
TOBACCO SUBSTATION AT WINDSOR. Report for 1933. P. J. Anderson, T. R. Swanback and O. E. Street. No. 359.
REPORT OF STATE ENTOMOLOGIST FOR 1933. W. E. Britton. No. 360.
CROSSED SWEET CORN. Donald F. Jones, W. Ralph Singleton. No. 361.
COMMERCIAL FEEDING STUFFS. Report on Inspection. E. M. Bailey. No. 362.
REPORT ON FOOD PRODUCTS AND DRUGS FOR 1933. E. M. Bailey. No. 363.
TOBACCO CULTURE IN CONNECTICUT. P. J. Anderson. No. 364.

CIRCULARS OF THE STATION

- Testing Vegetables for Connecticut. Results for 1933. L. C. Curtis. No. 94.
The Interpretation of Soil Tests. M. F. Morgan. No. 95.
Stewart's Bacterial Wilt on Sweet Corn. G. P. Clinton and W. Ralph Singleton. No. 96.
Fleas and Their Control. B. H. Walden. No. 97.
Late Blight of Tomatoes. A. A. Dunlap. No. 98.
Control of the Plum Curculio on Fruit Trees. Philip Garman. No. 99.
Substitutes for Lead Arsenate on Fruits and Vegetables in Connecticut. Philip Garman and Neely Turner. No. 100.
Control of Apple Maggot. Philip Garman. No. 101.
Potato Spraying. A. A. Dunlap and Neely Turner. No. 102.
Law and Regulations Concerning Inspection and Shipment of Nursery Stock in Connecticut. W. E. Britton. No. 103.
Quarantine Measures Restricting Shipments of Connecticut Plants, 1934. W. E. Britton. No. 104.
Regulations Concerning Transportation of Nursery Stock in the United States and Canada. W. E. Britton. No. 105.

JOURNAL PAPERS

- BLOCK, RICHARD J., and FARQUHAR, LUCILLE REED. Studies on vitamin G (B_2). I. Yeast and liver preparations as a source of vitamin G (B_2). *Jour. Biol. Chem.*, 103; 643-649. 1933.
BOTSFORD, R. C. Mosquito Control in Connecticut, 1933-1934. *Proc. 21st Annual Meeting, New Jersey Mosquito Extermination Association*, pp. 112-114, July 1934.
BRITTON, W. E. A Tropical Moth in Connecticut. *Entomological News*, Vol. XLV, p. 43, February 1934.
BRITTON, W. E. Additional Inspection of Nurseries on Account of the European Pine Shoot Moth. *Jour. Econ. Ent.*, Vol. 27, 572, June 1934.
BRITTON, W. E. The Corn Ear Worm and the European Corn Borer and Their Control. *Proc. 21st Annual Meeting, Conn. Veg. Growers' Assn.* p. 72, May 1934.
BRITTON, W. E. Experimental Work on Vegetable Insects in 1933. *Proc. 21st Annual Meeting, Conn. Veg. Growers' Assn.* p. 41, May 1934.
BRITTON, W. E. Injury to Trees by Squirrels. *Ninth Annual Shade Tree Conference*, p. 85, January 1934.
BRITTON, W. E. Miscellaneous Problems in Beekeeping. *Jour. Econ. Ent.*, Vol. 27, p. 596, June 1934.

- BRITTON, W. E. The Mosquito Problem of Connecticut and How to Solve It. 16 pp., 10 figs., (Published by State Department of Health). Revised edition. Hartford, January 1934.
- BRITTON, W. E. Need of Further Research on the European Corn Borer. Hearings before Sub-Committee of Committee on Appropriations, U. S. Senate on H. R. 8134, p. 67, March 1934.
- BRITTON, W. E. Report of Committee on Injurious Insects. Proc. 43rd Annual Meeting, Conn. Pomological Society, p. 110, April 1934.
- BRITTON, W. E. Report on Vegetable Insects in 1933. Proc. 21st Annual Meeting, Conn. Vegetable Growers' Assn. p. 42, May 1934.
- BRITTON, W. E. Some Insect Pests of Cultivated Plants. In Garden Guide, p. 411, (Published by A. T. De LaMare Co., New York). Revised edition. March 1934.
- CLARK, H. T., FOSTER, G. L., and VICKERY, H. B. Über die "neue Methode zur Darstellung von Aminen aus Aminosäuren" von Wada. Biochem. Ztschr., 272: 376-379. 1934.
- CLINTON, G. P. Plant Diseases in Connecticut. Proc. Connecticut Vegetable Growers' Assn. 1933: 46-50. 1934.
- CLINTON, G. P. Stewart's Bacterial Wilt of Sweet Corn. Proc. Connecticut Vegetable Growers' Assn. 1933: 69-72. 1934.
- DUNLAP, A. A. Spraying and Dusting of Tomatoes for Late Blight (*Phytophthora infestans*) Phytopathology, 24: 8. 1933.
- FISHER, H. J. Investigation of Methods for Determining Calcium Gluconate. Jour. Asso. Official Agri. Chemists. August 1934.
- FRIEND, R. B., and WEST, A. S., JR. Spray Experiments for the Control of the European Pine Shoot Moth. Jour. Econ. Ent., Vol. 27, p. 334, April 1934.
- FRIEND, R. B., and WILFORD, B. H. The Spruce Gall Aphid as a Forest Pest. Jour. Forestry, XXXI, No. 7, p. 816, November 1933.
- GARMAN, PHILIP. Studies on Control of the White Apple Leafhopper in Connecticut. Jour. Econ. Ent., Vol. 27, p. 361, April 1934.
- GARMAN, PHILIP. Report of Results of Tests with Lead Arsenate Substitutes. Proc. 43d Annual Meeting, Conn. Pomological Society, pp. 70-80, May 1934.
- HORST, KATHRYN, MENDEL, LAFAYETTE B., and BENEDICT, FRANCIS G. The Influence of Previous Exercise upon the Metabolism, the Rectal Temperature, and the Body Composition of the Rat. Jour. Nutrition, 7: 251-275. 1934.
- HORST, KATHRYN, MENDEL, LAFAYETTE B., and BENEDICT, FRANCIS G. The Effects of Some External Factors upon the Metabolism of the Rat. Jour. Nutrition, 7: 277-303. 1934.
- HORST, KATHRYN, MENDEL, LAFAYETTE B., and BENEDICT, FRANCIS G. The Influence of Previous Diet, Growth and Age upon the Basal Metabolism of the Rat. Jour. Nutrition, 8: 139-162. 1934.
- JACOBSON, H. G. M. Relative Influence of Nitrate and Ammonical Nitrogen upon Intake of Calcium by Tobacco Plants. Plant Physiology, 3: 340-342. 1933.
- JOHNSON, J. P. Corn Borer Clean-up Law, Hartford County Farm News, February 1934.
- JONES, DONALD F. Unisexual Maize Plants and Their Relation to Dioecism in Other Organisms. Proc. Nat'l. Academy of Sciences, Vol. 20, No. 1, p. 39, January, 1934.
- LUNT, H. A. Variations in Soil Texture as Revealed by Moisture Equivalent Determinations. Jour. Am. Soc. of Agron. 26: 713-715.
- MORGAN, M. F. Report on the Reaction Value of Acid Soils. Jour. Off. Agr. Chem. 17: 237, 1934.
- MORGAN, M. F. A System of Determining the Nutritive Status of the Soil by Microchemical Tests of a Single Soil Extract. Proc. of Northeastern Section, Am. Soc. of Agron., Boston, Mass., December 28, 1933: 11-17.
- MORGAN, M. F. Soil Testing as a Guide to Sound Soil Management. Am. Potato Journal, October 1933: 259-265.
- PLUMB, G. H., and HICOCK, H. W. Insect Control Work by the Civilian Conservation Corps Camps in Connecticut. Jour. Econ. Ent., Vol. 27, p. 344, April 1934.
- PUCHER, GEORGE W., VICKERY, HUBERT B., and LEAVENWORTH, CHARLES S. Determination of the Acids of Plant Tissue. III. Determination of Citric Acid. Ind. and Eng. Chem., Anal. Ed., 6: 190-192. 1934.

- PUCHER, GEORGE W., VICKERY, HUBERT B., and WAKEMAN, ALFRED J. Determination of the Acids of Plant Tissue. II. Total Organic Acids of Tobacco Leaf. *Ind. and Eng. Chem., Anal. Ed.* **6**: 140-143. 1934.
- PUCHER, GEORGE W., VICKERY, HUBERT B., and WAKEMAN, ALFRED J. Determination of the Acids of Plant Tissue. IV. A New Method for the Determination of Malic Acid. *Jour. Biol. Chem.* **105**: lxxviii-lxix. 1934.
- PUCHER, GEORGE W., VICKERY, HUBERT B., and WAKEMAN, ALFRED J. Determination of Malic Acid in Plant Tissue. Simultaneous Determination of Citric and Malic Acids. *Ind. and Eng. Chem., Anal. Ed.*, **6**: 288-291. 1934.
- SCHREAD, JOHN C., and GARMAN, PHILIP. Some Effects of Refrigeration on the Biology of Trichogramma in Artificial Breeding. *Jour. N. Y. Ent. Society*, XLII, pp. 263-283. Sept. 1934.
- STODDARD, E. M. Brown Rot Infection of Peaches in 1933 and Report of the Committee on Fruit Diseases. *Conn. Pom. Soc.* 1933: 41-44. 1934.
- STODDARD, E. M. Progress Report on Investigations on a New Peach Trouble. *Conn. Pom. Soc.* 1933: 115-117. 1934.
- VICKERY, HUBERT B. Report on Forms of Nitrogen in Plants. *Jour. Assoc. Off. Agr. Chem.* **16**: 473-476. 1933.
- VICKERY, HUBERT B. The Biochemistry of the Nitrogenous Constituents of the Green Plant. *Ann. Rev. Biochem.* **3**: 475-484. 1934.
- VICKERY, HUBERT B. Report on Forms of Nitrogen in Plants. *Jour. Assoc. Off. Agr. Chem.* **17**: 273-275. 1934.
- VICKERY, HUBERT B. What the Organic Chemist Can Do for Plant Physiology. *Plant Physiol.* **9**: 685-688. 1934.
- VICKERY, HUBERT B., and GORDON, WILLIAM G. Complex Compounds Formed by Certain Amino Acids in the Presence of Mercuric Chloride and Alkali. *Jour. Biol. Chem.* **103**: 543-547. 1933.
- VICKERY, HUBERT B., and PUCHER, GEORGE W. A Stillhead for Rapid Concentration in Vacuo. *Ind. and Eng. Chem., Anal. Ed.*, **6**: 372. 1934.
- VICKERY, HUBERT B., and WHITE, ABRAHAM. Proportion of Cystine Yielded by Hemoglobins of the Horse, Dog, and Sheep. *Proc. Soc. Expt. Biol. Med.*, **31**: 6-7. 1933.
- VICKERY, HUBERT B., and WHITE, ABRAHAM. The Basic Amino Acids of Casein. *Jour. Biol. Chem.* **103**: 413-415. 1933.
- YUDKIN, ARTHUR M., FARQUHAR, LUCILLE REED, and WAKEMAN, ALFRED J. Some Abnormalities in Rats Subsisting on Diets Poor in Mineral Nutrients. *Arch. Path.* **17**: 40-45. 1934.

All of which is respectfully submitted,

WILLIAM L. SLATE,
Director

INDEX TO ANNUAL REPORT

For the Year Ending October 31, 1934

Abbot sphinx moth	156, 164
<i>Acanthoscelides obtectus</i>	166
<i>Acholla multispinosa</i>	168
Acme Evans Co., feeds	316
Acme Guano Co., fertilizers	12, 23, 24, 29, 31, 32, 36
<i>Acroneuria abnormis</i>	167
<i>Acrosternum hilare</i>	158
<i>Acucephalus nervosus</i>	167
<i>Adelges abietis</i>	158, 171
<i>pinicorticis</i>	158
<i>strobilobius</i>	158
<i>Aegeria exitiosa</i>	154
<i>Agrilus bilineatus</i>	158
Agronomy department, see soil	
<i>Ailaus oculatus</i>	167
Albers Bros. Milling Co., feeds	316
Allen Co., E. T., feeds	317, 342
Allied Mills, Inc., feeds	317, 343, 353, 359, 376, 377, 378, 379
<i>Allocaënia pygmaea</i>	167
<i>Alsophila pometaria</i>	152, 154, 158, 213
<i>Alternaria tenuis</i>	131, 132, 133, 134
Aluminum, tested in soil	479
<i>Allypia octomaculata</i>	154
<i>Amaranthus retroflexus</i>	204
<i>Ambrosia artemisiifolia</i>	204
<i>trifida</i>	204
American Agricultural Chemical Co., fertilizers ..	12, 22, 23, 24, 27, 29, 31, 32, 36, 54
American cockroach	166, 255, 256
American Cyanamid Co., fertilizers	12, 20, 36
American Maize Products Co., feeds	317, 351
<i>Amphibolips inanis</i>	158
Anabasine, pure	238, 239
sulfate	229, 238, 239, 240
Analyses of Common Foods, compilation	529
Analytical Chemistry:	
commercial feeding stuffs, report on inspection	313
commercial fertilizers, report for 1934	5
common foods, analyses of	528
food and drugs, report for 1934	489
work reviewed in Director's report	69
<i>Anasa tristis</i>	156, 224
<i>Anastatus disparis</i>	196
Anchor Mills, feeds	317, 379
<i>Andricus seminator</i>	158
Anheuser-Busch, Inc., feeds	317
<i>Anisota rubicunda</i>	158, 249
<i>senatoria</i>	158
<i>Anomala orientalis</i>	153, 164, 165
<i>Anthrenus scrophulariae</i>	166
Ant-lion	168
<i>Auraphis roseus</i>	151, 155
<i>Apanteles congregatus</i>	168
sp.	232
<i>Apateia rubricoma</i>	159
Aphis lion	168

<i>Aphis pomi</i>	151, 155, 163
<i>rumicis</i>	238
<i>Aphrophora parallela</i>	159
Apiaries, inspection and statistics, 1934	81, 182
Apothecaries Hall Co., fertilizers	12, 13, 20, 22, 23, 24, 26, 29, 30, 31, 32, 36, 54
Apparatus and glassware for universal soil tests	463
Apple: maggot, winter breeding of	152, 156, 235, 236, 253
scab	210, 211, 212
seed chalcid	156
spray residue	491
sprays, tests of	210-212
Arborvitae leaf miner	159
soft scale	160
Arcady Farms Milling Co., feeds	318, 353, 359, 373, 379, 380
Archer-Daniels-Midland Co., feeds	318, 343
Argus tortoise beetle	157
<i>Argyresthia thuiella</i>	159
Armour Fertilizer Works, fertilizers	13, 23, 24, 29, 32, 38, 53, 54
Arsenical poisons	265
Arsenious acid, solution of, analysis	514
Ascogaster	231
<i>Ascogaster carpocapsae</i>	252
<i>quadridentata</i>	232, 252
Ashcraft-Wilkinson Co., fertilizers, feeds	12, 13, 22, 318, 342
Asiatic beetle	153, 164, 165
<i>Aspidiotus perniciosus</i>	151, 155
Associated Seed Growers, Inc., fertilizers	13, 20, 23, 24, 38
<i>Asterolecanium variolosum</i>	159
Atkins & Durbrow, Inc., fertilizers	13, 53, 54
<i>Attagemis piceus</i>	166
<i>Automeris io</i>	159
<i>Autoserica castanea</i>	153, 157, 164, 165
Babcock Glassware, etc., tested	528
Bacterial wilt	74
Bagworm	154, 163, 257, 258
E. W. Bailey & Co., feeds	318, 353, 359, 360, 373, 380, 381
The Baker Castor Oil Co. of Delaware, fertilizers	13, 22
Banding trees, for canker worms	272
for gypsy moths	643
Barberry webworm	152, 163
Barium fluosilicate	237, 246
treatment for bean beetle	441-451
Bark beetles and borers on elm	296-307
Barrett Co., fertilizers	13, 20
F. A. Bartlett Tree Expert Co., fertilizers	13, 38
<i>Bassus diversus</i>	232
Beacon Milling Co., Inc., feeds	318, 319, 353, 360, 373, 376, 381, 382
Bean: leafhopper	246
weevil	166
Bed bug	166
Beech: aphid	161
scale	154, 159, 220-224
control	224
Bentonite	235, 237
Berkshire Chemical Co., fertilizers	13, 22, 23, 24, 29, 30, 31, 32, 38
Beverages, carbonated, analysis	492
sweet cider, wine, analysis	492
<i>Bibio</i> sp.	167
Biochemistry, department of	
chemical investigations of the tobacco plant, bul. 374	557
review of work in 1934, Director's report	70, 72

Birch leaf mining sawfly	160
Bisbee Linseed Co., fertilizers, feeds	13, 29, 32, 319, 343
Bismuth, glycerite of, analysis	514
<i>Bittacomorpha clavipes</i>	167
Black blister beetle	164
carpet beetle	166
-headed pine sawfly	161
horse fly	168
vine weevil	153, 163, 164
Blackberry psyllid	156
Black Rock Milling Corp., feeds	319, 360, 361, 382
Blatchford Calf Meal Co., feeds	319, 376
<i>Blatta orientalis</i>	167
<i>Blissus leucopterus</i>	165
Blister rust control	82
Blue spruce gall aphid	160
Bordeaux mixture	236, 246, 247
on potatoes	74
The Borden Sales Co., Inc., feeds	319
Borers	
banded and red hardwood, on elm	306
occasional, on elm	307
Botany department	
Dr. Clinton's collections	94
work in 1934 reviewed	72-75
Box psyllid	164
-elder psyllid	162
<i>Brachyrhinus ovatus</i>	153, 159
<i>sulcatus</i>	153, 163, 164
Joseph Breck & Sons Corp., fertilizers	12, 54
Amos D. Bridge's Sons, Inc., feeds	319, 361
Broad-horned prionus	162
The Brode Corp., feeds	319, 342
L. Broder Grain Store, feeds	319, 361
Brown-Forman Dist. Co., feeds	319
Brown-tail moth, elm pest	279
control	281
C. Buckingham & Co., Inc., feeds	319, 320, 361, 382, 383
C. E. Buell, Inc., fertilizers, feeds	13, 20, 21, 22, 53, 54, 320, 409
Buffalo tree hopper	294
control	295
Bumble flower beetle	155, 160
C. W. Burckhalter, Inc., feeds	320, 409
Cabbage maggot	157
<i>Cacoecia argyrospila</i>	155
<i>rosaceans</i>	155
Cairo Meal and Cake Co., feeds	320
<i>Calaphis castaneae</i>	159
Calcium arsenate	235, 236, 246, 247
cyanide	255
<i>Callidium frigidum</i>	167
<i>Calligrapha philadelphica</i>	159
<i>Callosamia promethea</i>	163
<i>Calosoma calidum</i>	168
<i>sycophanta</i>	196
C. W. Campbell Co., feeds	320, 353, 354, 361, 373, 383
<i>Camponotus herculeanus</i>	167
Canker worms	
control	216
spraying	216
banding	217

fall	269, 271
outbreak of	213-220
spring	270
A. B. Caple Co., feeds	320, 354
Carbon disulfide	229
tetrachloride	251
Carpenter ant	167
worm	162
Carpet beetle	166
<i>Carpocapsa pomonella</i>	155
<i>Caryomyia caryaecola</i>	159
Case-bearing clothes moth	166
<i>Casnonia pennsylvanica</i>	168
Cassina	505, 506
<i>Catocala epione</i>	167
Cecropia moth	156
Cedar rust	212
Celery worm	158
Center Milk Products Co., feeds	320, 409
<i>Cerastipsocus venosus</i>	167
Chain-dotted geometer	163
Chapin & Co., feeds	320
Charts, calibration of, in soil tests	469
color in soil tests, plates I-VIII	384
<i>Chauliodes</i> sp.	167
Cheese, analysis	492
<i>Chelymorphe cassidea</i>	157
Chemical Investigations of the Tobacco Plant	
Part V, bul. 374	553-620
<i>Chenopodium album</i>	204
Chestnut blight	72
Chilean Nitrate Sales Corp., fertilizers	13, 20, 58
Chinch bug	165
Chinese praying mantid	152, 169
<i>Chionaspis cuonyni</i>	151, 163
<i>pinifoliae</i>	152, 159
Chlorinated soda, solution of, analysis	514, 515
<i>Chrysochus auratus</i>	167
<i>Chrysopa oculata</i>	168
Cicada killer	166
Cigarette beetle	166
<i>Cimex lectularius</i>	166
<i>Cincticornia pilulae</i>	159
sp.	159
<i>Cingilia catenaria</i>	163
<i>Citheronia regalis</i>	159
Clinton Co., feeds	320, 351
Herbert K. Clofine, feeds	320, 409
Clothes moths, control	252
Clover weevil	168
Clyde Renco Milling Corp., feeds	320
<i>Coccinella transversoguttata</i>	168
Codling moth	155, 211, 212, 235, 236
Coffee, etc., analysis	492
<i>Colaspis favaea</i>	155
<i>Coleophora laricella</i>	159
Coles Co., feeds	320, 361, 383
Collis Products Co., feeds	321, 409
Colorado potato beetle	157
Columbia Modified Soda, No. 200	523
Columbine borer	165
Commander-Larabee Corp., feeds	321, 344, 346, 349

Commercial Feeding Stuffs, inspection, 1934, bul. 370	310-318
Commercial Fertilizers, report for 1934, bul. 365	1-58
Common red spider	165
Common foods	529
analyses of	532
Community Service, Inc., feeds	321, 361, 383
Composite solution standards in soil testing	482
<i>Compsilura concinnata</i>	197
Confectionery, alcohol in	495
Conference of Connecticut Entomologists	170
G. E. Conkey Co., feeds	321, 383, 384
<i>Conistra indirecta</i>	163
Connecticut Fat Rendering and Fertilizer Corp., fertilizers, feeds ..	13, 29, 31, 321, 409
Connecticut State Entomologist, report for 1934, bul. 368	147-262
<i>Conotrachelus juglandis</i>	159
<i>nenuphar</i>	152, 155
Consolidated Products Co., feeds	321, 409
Consolidated Rendering Co., fertilizers, feeds	14, 20, 22, 23, 24, 26, 27, 29, 30, 31, 32, 40, 53, 54, 321, 410
Consumers Import Co., Inc., feeds	321, 410
Continental Distilling Corp., feeds	321
Copeland Flour Mills, Ltd., feeds	321, 344, 346
Copper	246
Copper-lime-calcium arsenate	441
Corethrine larvae in drinking water	167
Corn	
bacterial wilt on	74
crosses	
advanced generation	681
multiple	682
reciprocal	679
single	678
single plant	683
variety-inbred	683
whipple yellow sweet	671
white flint	663
evergreen sweet	667
testing and utilization of inbred strains	649
contents	651
Corn borer, see European	
Corn ear worm	152, 157
Corn Products Sales Co. of Corn Products Refining Co., feeds	321, 351
<i>Corydalis cornuta</i>	167
<i>Corythucha arcuata</i>	159
<i>Cosmopepla bimaculata</i>	164
<i>Cotalpa lanigera</i>	155
E. A. Cowee Co., feeds	321, 322, 361, 384
C. A. Cowles, Inc., feeds	322, 354, 361, 362, 384
Chas. M. Cox Co., feeds	322, 344
Crane fly	167
<i>Cremastus</i> sp.	232, 234
Crinkled flannel moth	155, 164
Crosby Milling Co., feeds	322, 385
Crown gall	181
Crown Mills, feeds	322
Cryolite	235, 236, 237
<i>Cryptococcus fagi</i>	154, 159, 220, 222, 224
<i>Cryptorhynchus lapathi</i>	159
<i>Ctenocephalus canis</i>	166
P. Cutler, Inc., feeds	322, 362, 373, 385
Cutworms	157, 165
Cyanogas	229

Cyclamen mite	165
<i>Cyllene caryae</i>	167
<i>robiniae</i>	159
Dairymen's League Cooperative Assoc., Inc., feeds	322, 411
<i>Danaus menippe</i>	152, 167, 251
Dark native elm bark beetle	298
<i>Dasyneura pyri</i>	155, 254
<i>Datana ministra</i>	155, 159
Davey Tree Expert Co., fertilizers	14, 40, 53, 54
R. G. Davis & Sons, Inc., feeds	322, 354, 362, 373, 385
Decatur Milling Co., Inc., feeds	322, 351
Definitions of fertilizers	7
Delaware Mills, Inc., feeds	322, 354, 362, 363, 373, 385
<i>Dendrothrips ornatus</i>	163
The Denver Alfalfa Milling & Products Co., feeds	323, 354
<i>Dermestes lardarius</i>	166
<i>nidum</i>	255
<i>peruvianus</i>	154, 166, 232
Derris dust	246
<i>Diabrotica duodecimpunctata</i>	164
<i>vittata</i>	157
<i>Diaphania nitidalis</i>	154, 157, 249
<i>Diapheromera femorata</i>	159
<i>Diaspis carueli</i>	159
<i>Dichelonyx elongata</i>	159
<i>Dichomeris marginellus</i>	152, 163
Dietrich & Gambrill, Inc., feeds	323, 363, 373, 376, 385
<i>Dilachnus</i> sp.	160
<i>Diocles molestae</i>	232
Director's report for 1934, bulletin 366	61-100
Disease survey	75, 93
Diseases	
tobacco leaf spots	117
vegetable	74
Dobson fly	167
Dogwood club gall	163
<i>Dorcus parallelus</i>	167
<i>Drepanaphis acerifoliae</i>	160
Drugs, report for 1934 and analyses	487-553
Dry ground fish	29
analyses of	30
Duluth-Superior Milling Co., feeds	323, 344
Arthur G. Dunn, feeds	323, 344, 345, 346
Dutch elm disease	i, 64, 65, 256, 257, 296
Eastern States Farmers' Exchange, fertilizers, feeds	
14, 20, 22, 23, 24, 25, 27, 28, 29, 30, 32, 40, 323, 354, 363, 374, 385, 386, 387	
Eastern tent caterpillar	152, 154, 156, 161
<i>Ecpantheria deflorata</i>	168
S. T. Edwards Co., Inc., feeds	323, 411
Eggs and egg products, analyzed	495, 496
Ed Eggert, fertilizers	14, 26
Eight-spotted forester	154
Eisemann & Co., Inc., feeds	323, 342
Elm, insect pests of	265
<i>aspidiotus</i>	293
bark beetle	297
bark weevil	299
common borer	300
flea beetle	287
lacebug	291

leaf aphid	289
leaf beetle	152, 160, 285
leaf miner	283
native bark beetle	298
span worm	273
twig aphid	289
woolly aphid	290
Elmore Milling Co., Inc., feeds	324, 355, 363, 374, 376, 387
<i>Elodia</i> sp.	232
Thomas W. Emerson Co., fertilizers	14
<i>Empfytus cinctus</i>	181
<i>Empoasca fabae</i>	157, 246, 247
Emporia Elevator and Feeding Co., fertilizers	14, 53, 55
Entomology, Director's report on	75
Entomologist's report	147
contents	149
Entomological features for 1934	151
<i>Epicauta cinerea</i> var. <i>marginata</i>	157
<i>pennsylvanica</i>	164
<i>Epilachna corrupta</i>	152, 157, 245
Epione, underwing moth	167
<i>Epitrix cucumeris</i>	157, 246
<i>Erax aestuans</i>	169
<i>Eriophyes eucricotes</i>	163
sp.	160
<i>Eriosoma americana</i>	160
<i>lanigera</i>	155
sp.	160
John W. Eshelman & Sons, feeds	324, 355, 363, 364, 374, 377, 387, 388
Ethyl nitrite	519, 520
<i>Eubadizon</i> sp.	234
<i>Euclea chloris</i>	168
<i>Eucosma gloriola</i>	160
<i>Eulia pinatubana</i>	160
<i>Euonymus radicans</i>	163
Euonymus scale	151, 154, 163
<i>Euphoria fulgida</i>	168
<i>inda</i>	155, 160
European corn borer	77, 153, 154, 158, 165, 245
control	198-206
damage to the corn crop	200
fall clean-up campaign	199
spring enforcement of the compulsory clean-up	198
weed survey	204
European	4, 296
elm bark beetle	151, 154, 294
elm scale	293
fruit lecanium	151, 154, 162, 171
pine shoot moth	156, 161
red mite	155
<i>Euthisanotia grata</i>	168
<i>Euxesta notata</i>	324, 351
Evans Milling Co., feeds	325, 349
Excelsior Milling Co., feeds	167
Eyed elater	325, 347
Fairchild Milling Co., feeds	325
The Fairmont Creamery Co., feeds	152, 154, 158, 213, 214, 215, 217, 269
Fall canker worm	153, 160, 281
webworm	160
False pine webworm	325, 353
Farmers Feed Co., feeds	

Farmers Flour & Grain Co., feeds	325, 388
Faroll Bros., feeds	325, 345
Fat and oils, analysis of	
butter	496, 497
olive oil	497, 498, 499, 500, 501
cottonseed oil	498
"salad oil"	497
Feed law, the	313
guides in enforcement	315
Feeding stuffs, terms used in analyses of	315
inspection of	341
<i>Fenusa pumila</i>	160
Fernando Valley Milling & Supply Co., feeds	325, 355, 388, 389
Fertilizer	
conservation by cover crops	90
experiments on tobacco	92
for sweet potatoes	88
market garden trials	88
Fertilizers, commercial	5
acid or alkaline effects	89
analyses of mixed	5, 18, 36, 58
definitions and interpretation of terms	7
inspection and analyses of, 1934	18
raw materials chiefly valuable for	
nitrogen	19
nitrogen and potash	27
nitrogen and phosphoric acid	29
phosphoric acid	23
potash	26
lime and acid reacting	90
miscellaneous fertilizers and other materials	53
check meals and fertilizers	58
liming materials	53
other materials	53
sheep manure and others	53
mixed fertilizers	34
special and home mixtures	35, 51
recent changes affecting control	6
registrations	12
statute and regulations regarding	5
Field Days, 1934	62
<i>Fiorinia japonica</i>	160
Fire brat	166
First National Stores, Inc., feeds	325, 389
Fish oil	210, 212, 246
Fleas	166
Flory Milling Co., Inc., feeds	325
Flotation sulfur	210, 211, 212, 235, 236, 237
Fluorine, insecticide for bean beetles	339, 451
residues on beans	451
Food and drug control	
Station's interest in	489
Food and Drug Products, annual report for 1934	487-553
Food, drugs, etc., miscellaneous samples examined	322, 487
Foods, common, analyzed	529
Ford Motor Co., fertilizers	14, 20
Forest soils, lysimeter data	90
moisture in	91
Forest tent caterpillar	153, 161, 249
Forestry, review of department work in Director's report	82-86
J. A. Forrest, feeds	325, 345, 347

Four-horned sphinx, elm pest	283
control	283
Four-lined plant bug	158
Freihofer Flour Mills, feeds	325
L. T. Frisbie Co., fertilizers	14, 29, 32
Fruit tree leaf roller	155
<i>Galerucella luteola</i>	152, 160
Garden millipede	157
J. B. Garland & Son, feeds	325, 364, 389
General Commodity Corp., feeds	329, 411
General Mills, Inc., feeds	326, 345, 347, 349
Genetics: review of department work in 1934, Director's report	84
Improvement of naturally cross-pollinated plants by selection in self-fertilized lines, bul. 376	651
Giant water bug	168
<i>Gillettea cooleyi</i>	160, 171
Gladiolus thrips	165
<i>Glischrochilus quadriguttatus</i>	168
Glue	237
<i>Glypta rufiscutellaris</i>	234
sp.	233
Golden glow aphid	164
tortoise beetle	164
Goldsmith beetle	155
<i>Gossyparia spuria</i>	151
Goulard and Olena, Inc., fertilizers	14, 42
Granary weevil	166
Grand Union Co., feeds	326, 390
D. H. Grandin Milling Co., feeds	326, 355, 365, 374, 377, 389
Grape phylloxera	156
plume moth	156
Grapevine tomato gall	155
<i>Graphium Ulmi</i>	1-8, 64, 65, 256, 257
<i>Grapholitha molesta</i>	152, 155
Grasselli Contact Insecticide, No. 1	229
Great Atlantic & Pacific Tea Co., feeds	326, 390
Green apple aphid	151, 155, 163
elm beetle	160
gold beetle	167
stink bug	158
striped maple worm	158, 249
Greenhouse whitefly	165
Ground fish, dry, analyses of	29, 30
Ground tobacco stems, analysis of	28
<i>Gynandropus hylacis</i>	169
Gypsum	229
Gypsy Moth, bul. 375	621-648
Gypsy moth	153, 154, 162, 221, 277
barrier zone	639
control	80, 188-197, 641-643
damage, destruction of trees	228, 229
description	633
distribution abroad and history in America	625-627
in Connecticut	627
imported parasites	636
life history	631
natural enemies	634
quarantine restrictions	639
<i>Hadronotus anasae</i>	228
<i>carinatifrons</i>	228

Hales & Hunter Co., feeds	326, 327, 390, 391
<i>Haltica ulmi</i>	160
Frank B. Ham & Co., Ltd., feeds	327, 345, 347
<i>Hamadryas antiopa</i>	160
William Hamilton & Son, Inc., feeds	327
Dwight Hamlin Co., feeds	327
Hardwood borers on elm	306
<i>Harpipteryx xylostella</i>	163
Hecker-H-O Co., Inc., feeds	327, 355
Hecker-Jones-Jewell Milling Div. of Standard Milling Co., feeds	327, 345, 347
The Dr. Heinz Co., fertilizers	14, 29, 33
<i>Heliothis obsoleta</i>	152, 157
<i>Heliothrips femoralis</i>	253
Hellgramite	167
<i>Hemaris thysbe</i>	163
<i>Hemerocampa leucostigma</i>	155
sp.	160
Hess Hair Milk	525
<i>Heterodera radicola</i>	163
Hickory bark beetle	162
borer	167
horned devil	159
seed gall	159
<i>Hippodamia convergens</i>	169
Hirst & Begley Lindseed Works, feeds	327, 343
Holly leaf miner	154, 164, 249
Paul M. Hubbard and Co., fertilizers	14, 42
The Hubinger Co., feeds	327, 351
Humphreys-Godwin Co., fertilizers, feeds	14, 22, 327, 342
Hydrated lime	210, 237
<i>Hylemyia brassicae</i>	157
<i>cilicrura</i>	157
<i>rubivora</i>	155
<i>Hylurgopinus rufipes</i>	5, 65, 256, 257, 298
<i>Hypera punctata</i>	168
<i>Hypermallus villosus</i>	152, 160
<i>Hyphantria cunea</i>	153, 160
Ice cream, etc., analysis of	502
<i>Ilex glabra</i>	164, 249
<i>opaca</i>	249
<i>Illinoia solanifolii</i>	157
Imperial Grain & Milling Co., feeds	327, 355
Imported willow leaf beetle	153, 162
Insect record for 1934	81, 154
Insects	
beneficial	168
flowers and greenhouse plants	164
fruit	154
household	166
infesting stored food products	166
infesting timber and wood products	167
ornamental shrubs and vines	163
shade and forest tree	158
soil and lawn	165
tobacco pests	93, 135-140
vegetable	156
Insect Pests of Elms in Conn., bul. 369	263-308
key	267
Inspection of apiaries	182-188
imported nursery stock	180-181
nurseries	171-180

International Agricultural Corp., fertilizers	14, 15
International Milling Co., feeds	327, 345, 347, 349
Io moth	159
Iodine, compound solution of, analysis	515
tincture of, analysis	516
Ipecac, fluid extract of, analysis	516
Iris weevil	164
<i>Itycorsia</i> sp.	160
Japanese beetle	153, 162, 165, 206-208, 287
inspection and certification	207
scouting	206
trapping	206
Japanese garden beetle	157, 164, 165
scale	161
weevil	153
Jersee Co., feeds	327, 391
<i>Julus hortensis</i>	157
June beetle	165
Juniper scale	159
webworm	152, 154, 163
The Kansas Flour Mills Corp., feeds	327
Kasco Mills, Inc., feeds	327, 365, 391
Kellogg Co., feeds	328, 351
Spencer Kellogg & Sons, Inc., fertilizers, feeds	15, 22, 328, 343
Kelloggs & Miller, Inc., feeds	328
<i>Kermes</i> , sp.	160
Kerosene emulsion	229
extract of pyrethrum	229, 230
-soap emulsion	224
Keystone Dairy Co., feeds	328, 411
H. H. King Flour Mills Co., feeds	328, 349
King Midas Mill Co., feeds	328
"Konate"	252
Kraft-Phenix Cheese Corp., feeds	328, 411
Chas. A. Krause Milling Co., feeds	328, 352
H. P. Kysor Feed & Grain Co., feeds	328, 355, 365, 391, 392
The Laden Bros. Co., Inc., feeds	328, 365, 392
Ladybeetle	169
convergent	169
five-spotted	168
<i>Lagoa crispata</i>	155, 164
John T. Lampman & Co., feeds	328
The Larabee Flour Mills Co., feeds	328, 345
Larch case bearer	159
woolly aphid	158
Larder beetle	166
Larger empty oak apple	158
The Larrowe Milling Co., feeds	328, 329, 342, 353, 365, 377, 392, 393
Late blight on tomatoes, control	74
"Larvex"	252
<i>Lasioderma serricorne</i>	166
<i>Lasioptera clavula</i>	163
vitis	155
<i>Lathridius liratus</i>	166
Lead arsenate	210, 211, 216, 235, 236, 237, 246
substitutes	76, 235-236
Leaf feeders, occasional, on elms	288
Leaf spot on tobacco	117
<i>Lecanium fletcheri</i>	160

Legal control measures, foods and drugs	489, 490, 491
Leopard moth, elm pest	302
control	304
<i>Lepidosaphes ulmi</i>	160, 163
<i>Leptinotarsa decemlineata</i>	157
Lesser European bark beetle	4, 256, 296
<i>Lethocerus americanus</i>	168
<i>Leucaspis japonica</i>	161
Libner Grain Co., Inc., feeds	329, 365, 374, 393
Library	94
Light-loving grapevine beetle	152, 156, 163, 165
Lima beans, control of bean beetle on	446
Lime	235, 236, 237, 246, 247
-sulfur	210, 212, 216, 224, 255
Limestone, analyses of	57
Lime-tree looper, pest of elms	272
control	273
The C. W. Lines Co., feeds	329, 365, 393
Linseed oil	210
<i>Lixus concavus</i>	57
Locust borer	159
Long Hill Feed Store, feeds	329, 365, 393
<i>Longistigma caryae</i>	161
Long-tailed mealybug	165
L. B. Lovitt & Co., fertilizers, feeds	15, 329, 342
Luna moth	168
Lunate long sting	169
<i>Lucilia sericata</i>	168
Luippold mineral water	523, 524
<i>Lycophotia margaritosa saucia</i>	164
<i>Lyctus linearis</i>	167
<i>Lygaeus kalmii</i>	163
<i>Lygus pratensis</i>	157
Lysimeter experiments	
data from forest soils	90
Macaroni, analysis of	505
<i>Macrobasis torsa</i>	161
Macrocentrus	231, 233
<i>Macrocentrus ancyliivorus</i>	232
<i>thoracicus</i>	232
<i>Macroductylus subspinosus</i>	155, 157
<i>Macrosiphum rudbeckiae</i>	164
Magnesia citrate, solution of, analysis	517, 518
Magnesium arsenate	246
in bean beetle control	441-452
Magnetic sulfur	210, 211, 212
dusting sulfur	253
Maine Fish Meal Co., feeds	329, 411
<i>Malacosoma americana</i>	152, 156, 161
<i>disstria</i>	153, 161, 249
<i>Mamestra picta</i>	157
E. Manchester & Sons, feeds	329, 347, 350, 365, 393
The Mann Bros. Co., feeds	329, 343
Maple bladder gall	162
spindle gall	161
Maple butter, analysis	523
Margined blister beetle	157
Maritime Milling Co., Inc., feeds	329, 330, 356, 366, 394, 395
Market garden fertilizer trials	88
E. J. Martenis Grain Co., feeds	330, 345
Maté (Paraguay tea)	505

Materials tested for poisons	527
Mating and productivity in rats	71
Meat products, analysis, etc.	
sausage	505, 507
hamburg steak	507
Meech & Stoddard, Inc., feeds	330, 366, 367, 374, 395
<i>Megarhyssa lunator</i>	169
<i>Melittia satyriniformis</i>	157
<i>Metritona bicolor</i>	164
Mexican bean beetle	77, 152, 157, 245, 246, 419
control experiments in Conn., bul. 371	419-452
date of planting	426-431
insecticides	439-445
poisonous	442, 444
poisonous residue on beans	449
non-poisonous	442, 444
spacing of plants	431-439
life history	425
Milk-weed bug	163
Milk and milk products, analysis, etc.	
cream	508, 509
evaporated milk	508
market milk	507
soft curd milk	507, 508
Millane Nurseries and Tree Experts Co., fertilizers	15, 42
The Miller Fertilizer Co., fertilizers	15, 23, 24, 42
Miner-Hillard Milling Co., feeds	330, 352
Minute brown scavenger beetle	166
Miscellaneous fertilizers and other materials	53
Miscellaneous foods and drugs, analyses	522, 523, 524, 525, 526, 527
Mite galls on elm leaves	161
Erineum caused by	160
on matrimony vine	163
Mites on <i>Euphorbia corollata</i>	164
Mixed fertilizers	34
analyses of	36
Monarch or milkweed butterfly	152, 167, 251
Monti-Van Iderstine, Inc., feeds	330, 411
<i>Monychus vulpeculus</i>	164
Geo. Q. Moon & Co., Inc., feeds	330, 331, 345, 347, 356, 367, 374, 395
Fred C. Morse & Son, feeds	331, 356, 367, 374, 395, 396
Jas. F. Morse & Co., feeds	331, 411
Mosaic	245, 247
Moseley & Motley Milling Co., feeds	331, 345, 347
Mosquito control	79, 208-209
Moth-proofing materials	78
Mount Carmel farm	
weather records	65
Mourning cloak butterfly	160
<i>Myrmeleon immaculatus</i>	168
Nantucket pine moth	162
Naphthalene	247
National Mineral Products Co., Ltd., feeds	331
Natural Guano Co., fertilizers	15, 53, 55
<i>Nectria coccinea</i>	223, 224
<i>Neoclytus acuminatus</i>	167
<i>Neodiprion lecontei</i>	161
pinetum	161
<i>Nepticula sericopeza</i>	161
<i>Nerice bidentata</i>	161
Niagara Falls Milling Co., feeds	331, 345, 347

Nicofume	229
Nicotine dust	229
pure	238, 239
solution	249, 253
sulfate	216, 229, 239, 240, 241
Nitrate of potash	27
analyses	28
Nitre, sweet spirits of, analysis	519
Nitrogen, in tobacco fertilizer	113
Nitrophoska, analysis of	48
Northwestern Consolidating Milling Div. of Standard Milling Co., feeds	331, 345, 347, 350
Norway maple leaf-stalk borer	161
Nurseries, inspection and statistics, 1934	80, 171-181
Nutrition investigations, review of work in 1934	71
N. V. Potash Export My., Inc., fertilizers	15, 27
Nu-Veg-Sal	522
Nux Vomica, tincture of, analysis	518, 519
Oak lacebug	159
mite	161
pea gall	161
pill gall	159
Oblique-banded leaf roller	155
Old Deerfield Fertilizer Co., Inc., fertilizers	15, 42
Olds & Whipple, Inc., fertilizers	15, 20, 23, 24, 26, 28, 29, 30, 31, 33, 42, 44
<i>Olethreutes hebesana</i>	164
<i>Omphalocera dentosa</i>	152, 163
Onion thrips	77, 247
<i>Opheltis</i> sp.	234
Orange-striped oak worm	158
Orchards, inspected	80
<i>Orgilus</i> sp.	232
Oriental cockroach	167
fruit moth	75, 152, 155, 233, 252
<i>Oryzaephilus surinamensis</i>	166
S. V. Osborn Estate, feeds	331, 356, 396
<i>Osmodermia scabra</i>	165
Outpost Nurseries, Inc., fertilizers	15, 44
<i>Oxyptilus periscelidactylus</i>	156
Oyster-shell scale	160, 163, 171, 292
<i>Pachystethus lucicola</i>	152, 156, 163, 165
Pacific Bone, Coal and Fertilizer Co., feeds	331
<i>Paleacrita vernata</i>	215
<i>Panchlora cubensis</i>	168
<i>Papaipema nitela</i>	158, 165
<i>purpurifascia</i>	165
<i>Papilio glaucus turnus</i>	161
<i>polyxenes</i>	158
<i>troilus</i>	164
Paraguay tea — see Maté	
Parasite of <i>Olethreutes hebesana</i>	169
of the Oriental fruit moth	231-234
Gypsy moth	636
<i>Paratetranychus bicolor</i>	161
<i>pilosus</i>	156, 161
<i>ununguis</i>	152, 161
<i>Paria canellus</i> var. <i>atterrimus</i>	161
var. <i>quadriguttatus</i>	161
Park & Pollard Co., feeds	331, 332, 350, 356, 357, 367, 368, 374, 397
The Patent Cereals Co., feeds	332, 352

Peach borer	154
soils of Connecticut	87
trees, "X" disease on	72
Pear leaf midge	155, 254, 255
psylla	152, 156
Pearl wood-nymph	155
Pecos Valley Alfalfa Mill Co., feeds	332, 357
Pedigreed Seed Co., Inc., fertilizers	15
<i>Pegomyia hyoscyami</i>	158
<i>Pelidnota punctata</i>	156
Penetrol extract pyrethrum	240
Penick & Ford, Ltd., Inc., feeds	332, 351
Pesticide	229
Peppers, new hybrids	85
<i>Periplaneta americana</i>	166, 255
<i>Perisierola angulata</i>	231, 232
Perkins Oil Co., fertilizer	12
<i>Perlesta placida</i>	168
<i>Phanerotoma grapholithae</i>	232
<i>Philonix pezomachoides</i>	161
sp.	161
<i>Phlegethontius quinquemaculata</i>	158
Phosphorus, fertilizer tests on tobacco	108-112
<i>Phyllaphis fagi</i>	161
<i>Phyllocoptes aceris-crumena</i>	161
<i>quadripes</i>	162
<i>Phyllophaga</i> sp.	165
<i>Phylloxera vitifoliae</i>	156
<i>Phymatodes variabilis</i>	167
<i>varius</i>	167
<i>Physokermes piceae</i>	162
<i>Phytomyza ilicis</i>	154, 164, 249
Pickle worm	154, 157, 249
Pickles, sweet, analyses, etc.	509
Pigeon tremex, elm pest	305
control	306
Pillbug	168
Pillsbury Flour Mills Co., feeds	332, 345, 348, 349, 350
Maurice Pincoffs Co., fertilizers, feeds	12, 15, 22, 332, 342
Pine bark aphid	158
blister rust	171
leaf scale	171
needle scale	152, 154, 159, 171
spittle bug	159
tube moth	160
Pine shoot moth	83
Pine weevil damage	83
<i>Pissodes approximatus</i>	162
<i>strobi</i>	153, 162
Pit-making oak scale	159
Pittsburgh Plate Glass Co., feeds	332
<i>Plagioderma versicolora</i>	153, 162
Plant Breeding	
lima beans	86
new Cuban shade tobacco	93
new squash	85
pepper hybrids	85
strawberries	86
testing and utilization of inbred strains of corn, bul. 376	650-678
contents	651
effect of deleterious factor on yield	686
parental characters in hybrids	661

performance of inbred strains when crossed	657
producing inbred strains	685
seed production	686
summary	689
testing inbreds	677
tomatoes	86
Plantspur Products Co., fertilizer	15
The Frank S. Platt Co., fertilizers, feeds	15, 44, 332, 397
Plum curculio	152, 155, 210, 235, 236
<i>Podisus maculiventris</i>	169
<i>Poecilocapsus lineatus</i>	158
<i>Popillia japonica</i>	153, 162, 165
Poplar and willow curculio	159
borer	162
canker	171
<i>Porcello scaber</i>	168
<i>Porthetria dispar</i>	153, 162
Post treatment	
plant established	84
tobacco poles	84
Postum Co., Inc., feeds	332, 352
Potash	
muriate of, analyses of	26, 27
soap	229, 230
sulfate of, analyses of	26, 27
Potato flea beetle	79
Potatoes, bordeaux mixture on	74
Potato aphid	157
flea beetle	157, 246
leafhopper	157
Pratt Food Co., feeds	333, 352, 357, 368, 397, 398
Precipitated bone	23
analyses of	25
Premier Poultry Manure Co., fertilizers	15, 53, 55
<i>Prionoxystus robiniae</i>	162
<i>Prionus laticollis</i>	162
<i>Pristomerus vulnerator</i>	232
<i>Prociophilus tessellatus</i>	162
Pro-Ker Milk, No. 10	525
Promethea moth	163
Protein chemistry, review of work in 1934	70
<i>Pseudocneorrhinus setosus</i>	153, 164
<i>Pseudococcus adonidum</i>	165
<i>Pseudolucanus capreolus</i>	166
<i>Psyllia buxi</i>	164
negundinis	162
pyricola	152, 156
<i>Pterodella pedicularis</i>	166
H. C. Puffer Co., feeds	333, 368, 398
Pulverized Manure Co., fertilizers	15, 53, 55
<i>Pyrausta nubilalis</i>	153, 158, 165
Pyrethrum	229, 230, 231, 240, 246, 253
treatment for bean beetles	443-451
<i>Pyrophila pyramidioides</i>	168
The Quaker Oats Co., feeds	333, 352, 357, 368, 375, 377, 398, 399
Radio, spray and weather reports	75
Ralston Purina Co., feeds	333, 334, 351, 357, 369, 375, 377, 399, 400
Raspberry cane maggot	155
John Reardon & Sons Co., feeds	334, 411
Red-headed pine sawfly	161


-humped caterpillar	156
Registrations of commercial fertilizers	12
<i>Reticulitermes flavipes</i>	167, 241
<i>Rhagoletis pomonella</i>	152, 156, 253
Rhubarb curculio	157
<i>Rhyacionia buoliana</i>	151, 162
<i>comstockiana</i>	162
<i>frustrana</i>	162
Robin Hood Mills, Ltd., feeds	334, 345, 348, 349
The Rogers & Hubbard Co., fertilizers	16, 23, 25, 29, 30, 33, 44, 46, 53, 55
Ronck & Bevis Co., feeds	334, 411
Root knot eelworms	163
<i>Rosa manetti</i>	180
<i>multiflora</i>	180
Rose chafer	155, 157
Rosy apple aphid	151, 155
Rotecide	229
Rotenone	246, 252
Rotex Milling Co., feeds	334, 400
H. M. Rubin Co., Inc., feeds	335, 411, 412
Russell-Miller Milling Co., feeds	335, 346, 348, 350
F. Rynveld and Sons, fertilizers	16
Saddle-back caterpillar	156
<i>Samia cecropia</i>	156
San José scale	151, 155, 171
San-O-Clean	523
<i>Saperda calcarata</i>	162
Satin moth	162
Sawfly larvae on arborvitae	162
on dogwood and hawthorn	164
on various kinds of pine	162
on white oak	162
Saw-toothed grain beetle	166
Scale, on elms	
<i>aspidiotus</i>	293
elm scurfy	293
European elm scale	294
European fruit lecanium	293
oyster-shell	292
<i>Schizura concinna</i>	156
<i>Scolytus multistriatus</i>	4, 256, 257, 296
<i>quadrispinosus</i>	162
<i>rugulosus</i>	156
O. M. Scott and Sons Co., fertilizers	16, 46
Seed corn maggot	157
Seed testing service	75
<i>Serica</i> sp.	166
Sewerage Commission of the City of Milwaukee, fertilizers	16, 46
Seymour Grain & Coal Co., feeds	335, 357, 369, 375, 401
Sheep manure and other materials, analyses of	53, 54
Sheffield Farms Co., Inc., feeds	335, 412
Shellabarger Grain Products Co., feeds	335
Sherwin-Williams Co., feeds	335, 343
M. L. Shoemaker & Co., Inc., fertilizers	16, 46
Shot-hole borer	156
<i>Sibine stimulea</i>	156
<i>Sinea diodema</i>	169
<i>Sira nigromaculata</i>	168, 253
Siroil	525
<i>Sitophilus granaria</i>	166
Slim	524

Small European elm bark beetle	4, 297
Small German cockroach	256
Snow white linden moth, pest of elms	273
control	274
Sodium aluminum silicofluoride	252
fluoride	253
oleate	238, 239
Soil, see also universal testing system	
and land cover inventory	86
extracting solution, choice of	457
in Connecticut peach orchards	87
reaction	470
sample for universal test	459
testing	87, 461, 464
universal system	457-484
Soils department, review of work in 1934, Director's report	86-91
Solutions and reagents, preparation of	461-463
universal soil extracting solution	461
Sooty blotch	212
Ike Sovitsky, feeds	335, 369, 375, 401
Sowbug	168
Special and home mixtures, fertilizers, analyses of	51
Sperry Flour Co., feeds	335, 350
<i>Sphecius speciosus</i>	166
<i>Sphecodina abbotii</i>	156, 164
Sphinx, see four-horned sphinx	
Spinach leaf miner	158
Spiny elm caterpillar	160, 274
control	274
Spotted grapevine beetle	156
Spray, residue	491
oil	80
Sprays for :	
bean beetle control	439, 451
canker worms	216
elm pests, see elm	
Gypsy moth	80, 188-197, 641
leaf eating insects	265
potatoes	74
Spring canker worm	213, 215
Spruce gall aphid	158, 171
gall scale	162
mite	152, 154, 161
Squash borer	157
bug	156, 224-231
control	227-231
Stag beetle	166
A. E. Staley Mfg. Co., feeds	335, 344, 351
Stalk borer	158, 165
Standard Wholesale Phosphate and Acid Works, Inc., fertilizers	
16, 20, 23, 25, 27, 29, 33, 46, 53, 56	
John T. Stanley Co., Inc., feeds	335
St. Albans Grain Co., feeds ...	336, 348, 350, 352, 353, 358, 370, 375, 377, 401, 402, 403
State Water Commission, examination of samples	527
D. A. Stickell & Sons, Inc., feeds	335, 336, 358, 370, 403
<i>Stilpnotia salicis</i>	162
St. Laurence Flour Mills Co., Ltd., feeds	336
Strawberry	
new varieties	86
root weevil	153, 154, 159
String beans, bean beetle control on	426-450
Striped cucumber beetle	157


<i>Sturmia scutellata</i>	197
Chas. M. Struven Co., feeds	336, 412
Sucking insects on elms	289, 296
Sulphuric acid, dilute, analysis	520
Summers Fertilizer Co., Inc., fertilizers	16, 23, 25, 29, 33, 46, 53, 56
Swallow-tail butterfly, green	164
tiger	161
Swift and Co. Fertilizer Works, fertilizers	16, 29, 33, 46
Synthetic cryolite	235
Synthetic Nitrogen Products Corp., fertilizers	16, 20, 48
<i>Syntomaspis druparum</i>	156
Syracuse Milling Co., feeds	337, 348, 358, 371, 375, 403, 404
<i>Tabanus atratus</i>	168
<i>Taeniothrips gladioli</i>	165
Tankage	29
analyses of	31
Tarnished plant bug	157
<i>Tarsonemus pallidus</i>	165
Tennessee Corp., fertilizers	16, 48
<i>Tenodera sinensis</i>	152, 169
Termite	167
damage	78, 241-245
<i>Tetralopha robustella</i>	162
<i>Tetranychus telarius</i>	165
<i>Thecodiplosis liriiodendri</i>	162
<i>Thermobia domestica</i>	166
I. P. Thomas & Son Co., fertilizers	16, 23, 25, 29, 31, 48
Thomaston Supply Co., feeds	337, 371, 404
Thrips on dahlia	165
on <i>Dianthus</i>	165
on Japanese iris	165
<i>Thrips tabaci</i>	247, 253
<i>Thyridopteryx ephemeraeformis</i>	154, 163, 257
<i>Tibicen canicularis</i>	168
<i>Tinea pellionella</i>	166
<i>Tineola biselliella</i>	252
Tioga-Empire Feed Mills, Inc., feeds	337, 349, 358, 371, 375, 377, 404, 405
Tobacco By-Products and Chemical Corp., fertilizers	16, 28
Tobacco, chemical investigations of plant	70, 557
analytical methods	563
contents of bulletin 374	555
description of collection	561
distribution of constituents	595
growth of plant in chemical terms	573
summary	604
work of Smirnov	602
Tobacco worm	158
Tobacco	
angular leafspot	120
blackfire	120
brown leaf spot and white speck	128
John Williams broadleaf spot	126
mosaic rust or fleck	126
new Havana seed	92
ringspot	120
wildfire	118
Tobacco insects, 1934	135
flea beetle control	137
hornworm control	139
thrips control	139
wireworm control	140

Tobacco fertilizer tests	
nitrogen	113-117
phosphorus	108-112
Tobacco Substation, annual report, bul. 367	103-145
insects in 1934	135
leaf spots	117
phosphorus in the fertilizer	108
review of work in Director's report	91
seed from one broadleaf plant	144
shade cloth and light intensity	143
tests on quantity of fertilizer	
nitrogen	113
<i>Tolyte velleda</i>	156
Tomatoes, breeding experiments	86
<i>Tomostethus bardus</i>	163
<i>Tortrix quercifolia</i>	163
<i>Toumeyella lirioidendri</i>	152, 163
Traders Feed & Grain Co., Inc., feeds	337, 346, 349
Transit Milling Co., feeds	337, 342
Tree hopper, see Buffalo	
"Tree Tanglefoot"	218, 220
<i>Trialeurodes vaporariorum</i>	165
Trichogramma	231, 232
<i>Trichopoda pennipes</i>	227
Jacob Trinley & Sons, Inc., feeds	337, 338, 358, 371, 405
<i>Trioza tripunctata</i>	156
<i>Triphleps insidiosus</i>	169
<i>Tropaea luna</i>	168
Truempy, Fasey and Besthoff, Inc., fertilizers	16, 29, 33
Tulip spot gall	162
tree scale	152, 163
Turpentine, report on	521
Twig girdlers, elm pests	304
Twelve-spotted cucumber beetle	164
Twig pruner	152, 154, 160
Two-lined chestnut borer	158
<i>Typhlocyba pomaria</i>	156
United Cooperative Farmers, Inc., feeds	338, 371, 405
United Mills Co., Inc., feeds	338
Union Starch & Refining Co., feeds	338, 351
Unity Feeds, Inc., feeds	338, 358, 371, 375, 377, 405, 406
Universal Soil Testing System, bul. 372	457-484
calibration of charts	469
choice of extracting solution	457
composite solution standards	482
interpretation of tests	470
preparation of solutions and reagents	461
procedure for conducting tests	464
plant tests	481
soil sample	459
soil reaction	470
supplemental tests	467
Upper Hudson Rye Flour Mills, Inc., feeds	338, 352
Urbana Mills Co., feeds	338, 412
Urea, analyses of	20
Van Iderstine Co., feeds	339
Variegated cutworm	164
Vegetables, studies in diseases	74
Victor Flour Mills, Inc., feeds	339, 346, 349, 350

Vinegar, analyses, etc.	514
Virginia-Carolina Chemical Corp., fertilizers	17, 48
S. A. Wald & Co., Inc., feeds	339, 342, 343
Walker-Gordon Laboratory Co., Inc., fertilizers	17, 53, 56
Hiram Walker & Sons, Inc., feeds	339
Walkingstick	159
Walnut weevil or curculio	159
Ward Dry Milk Co., feeds	339, 412
Watertown Cooperative Assoc., feeds	339, 407
Wayne County Grangers Feed Corp., feeds	339, 359, 372, 407
Weather records at Mount Carmel farm	65-68
Webbing clothes moth	252
F. S. Wertz & Son, feeds	339, 359
West-Nesbitt, Inc., feeds	339, 359, 372, 407
Wettable sulfur	237
White ants	167
White apple leafhopper	80
White-marked tussock moth, elm pest	274
control	275
White pine tip moth	160
White pine weevil	82, 153, 162, 171, 247, 248
Wilson's Original Compound Wa-Hoo Bitters	524
Wireworms	158, 166
Witch hazel, analysis	521, 522
Wolf's Feed Store, feeds	339, 372, 407
Thomas Wood & Sons, Inc., fertilizers	17
F. H. Woodruff & Sons, fertilizers	17, 48
Wool sower	158
Woolly alder aphid	162
apple aphid	155
Worcester Rendering Co., feeds	339, 412
"X" disease on peaches	72
<i>Xylotrechus colonus</i>	163
Yantic Grain & Products Co., feeds	340, 349, 350, 359, 372, 373, 375, 407, 408, 409
Yellow-necked caterpillar	155, 159
Yellows	245
Zebra caterpillar	157
Zinc sulfate	235, 236, 237
















University of
Connecticut
Libraries



39153028906586

